

**Global Meteorological Drought: A Synthesis of Current Understanding with a  
Focus on SST Drivers of Precipitation Deficits**

S. Schubert<sup>\*1</sup>, R. Stewart<sup>2</sup>, H. Wang<sup>1,3</sup>, M. Barlow<sup>4</sup>, H. Berbery<sup>5</sup>, W. Cai<sup>6</sup>, M. Hoerling<sup>7</sup>,  
K. Kanikicharla<sup>8</sup>, R. Koster<sup>1</sup>, B. Lyon<sup>9</sup>, A. Mariotti<sup>10</sup>, C. R. Mechoso<sup>11</sup>, O. Müller<sup>12</sup>, B.  
Rodriguez-Fonseca<sup>13</sup>, R. Seager<sup>14</sup>, S.I. Seneviratne<sup>15</sup>, L. Zhang<sup>16</sup>, T. Zhou<sup>16</sup>

<sup>1</sup>Global Modeling and Assimilation Office, NASA GSFC, Greenbelt, MD, USA

<sup>2</sup>University of Manitoba, Department of Environment and Geography, Winnipeg, Canada

<sup>3</sup>Science Systems and Applications, Inc., Lanham, MD, USA

<sup>4</sup>University of Massachusetts Lowell, Lowell, MA, USA

<sup>5</sup>Earth System Science Interdisciplinary Center/Cooperative Institute for Climate and  
Satellites, University of MD, College Park, MD, USA

<sup>6</sup>CSIRO Marine and Atmospheric Research, Aspendale, Victoria, Australia

<sup>7</sup>NOAA Earth System Research Laboratory, Boulder, CO, USA

<sup>8</sup>Indian Institute of Tropical Meteorology, Pune, India

<sup>9</sup>International Research Institute for Climate and Society, The Earth Institute, Columbia  
University, Palisades, NY, USA

<sup>10</sup>NOAA/OAR Climate Program Office, Silver Spring, MD, USA

<sup>11</sup>Department of Atmospheric and Oceanic Sciences, University of California, Los  
Angeles, CA, USA

<sup>12</sup>CEVARCAM, Facultad de Ingeniería y Ciencias Hidricas 5, Universidad Nacional del  
Litoral and CONICET, Santa Fe, Argentina

<sup>13</sup>Departamento de Física de la Tierra, Astronomía y Astrofísica-I, Facultad de CC.  
Físicas, Madrid, Spain.

<sup>14</sup>Lamont Doherty Earth Observatory of Columbia University, Palisades, NY, USA

<sup>15</sup>Institute for Atmospheric and Climate Science, ETH, Zürich, Switzerland

<sup>16</sup>LASG, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

Revision Submitted to

Journal of Climate GDIS Special Collection

19 January 2016

\*Corresponding author address: Siegfried Schubert, 8800 Greenbelt Rd.,  
NASA/GSFC, Greenbelt, MD 20771. E-mail: [siegfried.d.schubert@nasa.gov](mailto:siegfried.d.schubert@nasa.gov)

## ABSTRACT

Drought affects virtually every region of the world, and potential shifts in its character in a changing climate are a major concern. This article presents a synthesis of current understanding of meteorological drought, with a focus on the large-scale controls on precipitation afforded by sea surface temperature (SST anomalies), land surface feedbacks, and radiative forcings. The synthesis is primarily based on regionally-focused articles submitted to the Global Drought Information System (GDIS) collection together with new results from a suite of atmospheric general circulation model experiments intended to integrate those studies into a coherent view of drought worldwide. On interannual time scales, the preeminence of ENSO as a driver of meteorological drought throughout much of the Americas, eastern Asia, Australia and the Maritime Continent is now well established, whereas in other regions (e.g., Europe, Africa and India), the response to ENSO is more ephemeral or non-existent. Northern Eurasia, central Europe as well as central and eastern Canada stand out as regions with little SST-forced impacts on precipitation on interannual time scales. Decadal changes in SST appear to be a major factor in the occurrence of long-term drought, as highlighted by apparent impacts on precipitation of the late 1990s “climate shifts” in the Pacific and Atlantic SST. Key remaining research challenges include: (i) better quantification of unforced and forced atmospheric variability as well as land/atmosphere feedbacks, (ii) better understanding of the physical basis for the leading modes of climate variability and their predictability, and (iii) quantification of the relative contributions of internal decadal SST variability and forced climate change to long-term drought.

## 1. Introduction

Drought, which can occur in almost any region of the world, is one of the most destructive natural hazards faced by society. Some of the most dire concerns associated with climate change are associated with possible changes in drought frequency and severity, although regional drought projections often show large uncertainties (e.g., Seneviratne et al. 2012a; Orłowsky and Seneviratne 2013).

A substantial amount of research and operational effort has been devoted to drought. Many drought research studies have focused on particular regions or selected events, whereas others have examined the global distribution of droughts, their forcing factors, and their predictability. Efforts in operational environments now routinely assess current and future drought conditions over a variety of temporal and spatial scales. This broad range of activities, as well as many drought impact studies, suggests a need to document our collective understanding of and capabilities to predict drought. A synthesis of current understanding would help people everywhere benefit as much as possible from existing research and operational capabilities, through, for example, improved decision support and drought mitigation.

The Global Drought Information System (GDIS)<sup>1</sup> addresses these issues. The overall goal of GDIS is to provide coordinated information, monitoring, and prediction of drought

---

<sup>1</sup> The GDIS was developed as one of the key recommendations of a WCRP workshop on “Drought Predictability and Prediction in a Changing Climate: Assessing Current Knowledge and Capabilities, User Requirements and Research Priorities,” that was held on 2-4 March 2011 in Barcelona, Spain ([http://drought.wcrp-climate.org/workshop/ICPO\\_161\\_WCRP\\_Report.pdf](http://drought.wcrp-climate.org/workshop/ICPO_161_WCRP_Report.pdf)). The capabilities and requirements of the GDIS were further scoped out at a second workshop held in Frascati, Italy on 11-13 April 2012: (<http://www.clivar.org/organization/extremes/resources/dig>). A third workshop, “An

78 worldwide in a user-friendly manner. One of GDIS's objectives is to assess our current  
79 understanding of drought and our ability to predict it, thereby identifying research gaps.  
80 The present special collection of regionally-focused summary articles stems from this  
81 component of GDIS. Each article can stand on its own as an important contribution to  
82 drought research.

83  
84 It is also important, of course, to place these summary articles into context and to  
85 synthesize some of their findings. This is the goal of the present overview article. To  
86 make it tractable, we focus primarily on understanding the role of SST in driving  
87 meteorological drought, although some attention is also paid to other drivers as well as  
88 temperature anomalies. Furthermore, some of our discussion will focus more generally  
89 on seasonal-scale precipitation deficits, given that meteorological droughts can be  
90 considered extreme manifestations of such deficits – indeed, the level of deficit required  
91 to define a meteorological drought is not set in stone. In discussing such deficits  
92 generally, we make the implicit assumption that if a given set of conditions (as identified  
93 in this paper) leads to a seasonal precipitation deficit, then a more extreme version of  
94 these conditions would lead to a more extreme deficit and thus potentially a true  
95 meteorological drought. That is, we make the assumption that uncovering the sources of  
96 precipitation deficits on seasonal timescales is tantamount to uncovering the sources (if  
97 conditions therein were stronger) of meteorological drought.

---

International Global Drought Information System Workshop: Next Steps", was held in Pasadena on the Caltech campus 10-13 December 2014, and focused on reviewing the GDIS special collection papers, and developing the necessary next steps required for moving forward with an experimental real time global drought monitoring and prediction system (<http://www.wcrp-climate.org/gdis-wkshp-2014-about>).

In this paper we do not address how meteorological drought propagates to agricultural or hydrological droughts, or how soil moisture feedbacks, temperature changes, or human water use act to maintain or even amplify the different types of drought, though these issues are addressed to varying degrees in the articles of this GDIS collection.

Such a focus does not come without limitations; for example, the impact of long-term evapotranspiration changes induced by temperature and radiation changes (e.g., from climate change) may turn out to be as important as (if not more important than) precipitation changes in some regions in producing soil moisture and streamflow deficits at longer time scales (e.g., Cook et al. 2014; Cook et al. 2015). For example, Cook et al. (2014) used CMIP5-driven Palmer Drought Severity Index (PDSI) and Standardized Precipitation-Evapotranspiration Index (SPEI, Vicente-Serrano et al. 2010) drought estimates to show that, while robust regional changes in hydroclimate are primarily organized around regional changes in precipitation, increased potential evapotranspiration (PET) computed with the Penman-Monteith approach nearly doubles the percentage of global land area projected to experience significant drying based on these indices by the end of the 21st-century. Nevertheless, Sheffield et al. (2012), in addressing whether such impacts of increased PET are already evident in recent observationally-driven PDSI trends, found that global drought has changed little over the past 60 years (see also Seager and Hoerling 2014 for a discussion of regional differences), indicating that the focus here on precipitation deficits allows us to address much of the overall drought problem in current climate. Table 1 provides further

evidence that such a focus is justified, showing for our selected regions (see Section 3) generally high correlations between precipitation and either the PDSI or soil moisture.

We start by providing an overall scientific context for drought through an examination of the global drivers of precipitation and temperature changes on interannual and decadal time scales (Section 2). Next, we relate these and other factors to drought in different regions, highlighting implications for predictability and prediction (Section 3). Section 4 provides some concluding remarks.

## **2. Overview of Large Scale Factors**

Here we review the large-scale or the ultimate (as opposed to the proximate) causes of meteorological drought - the processes responsible for the long-term disruptions of local and regional precipitation-producing phenomena. These processes often act over large distances via various large-scale atmospheric motions such as the Hadley and Walker circulations, Rossby waves, and other atmospheric teleconnection patterns. The forcing for some of these large-scale motions is known to include sea surface temperature anomalies (SST), land (especially soil moisture) feedbacks, aerosols, and other natural and anthropogenic changes in radiative forcing such as those associated with global warming. These forcings are important because they may provide some degree of drought predictability (e.g., Smith et al. 2012). It must be kept in mind, however, that there is a substantial unforced (i.e., driven by processes internal to the atmosphere) element to the large-scale motions that significantly limits our ability to predict drought at the longer leads.

144

145 The various papers in the GDIS drought special collection assess, from a regional  
146 perspective, the global processes associated with meteorological drought. We summarize  
147 these findings here, while also providing some additional background on climate change  
148 aspects and regarding meteorological drought on the European continent. In addition, to  
149 provide a global framework for our discussion and synthesis, we include a model-based  
150 assessment of the dominant large-scale forcing of meteorological drought on seasonal  
151 and longer time scales – the response of the atmosphere to SST anomalies (e.g., Hoerling  
152 and Kumar 2003; Schubert et al. 2004; Seager et al. 2005). This assessment is based on  
153 AMIP-style simulations using prescribed SSTs (see Appendix A), with 5 different state-  
154 of-the-art global climate models; results are presented as combined (rather than  
155 individual model) statistics. We provide the model results in each subsection partly to  
156 assess their consistency with the findings of the individual special issue GDIS papers.  
157 The model results also provide insight into the spatial coherence and seasonality of the  
158 forced responses. In examining these results, we must keep in mind that their usefulness  
159 may be limited by model deficiencies and by the limitations imposed by employing SST-  
160 prescribed, integrations.

161

162 The link between drought and remote SST anomalies is complicated by the fact that there  
163 are different definitions of drought reflecting a wide range of societal (e.g., health, water  
164 quality, political), economic (e.g., agriculture, water supply, transportation, recreation)  
165 and ecosystem (e.g., fish, wildlife, wetlands, biodiversity, forest fires) impacts<sup>2</sup>. All of  
166 these definitions are important. Nevertheless, we focus here on the primary

---

<sup>2</sup> More on impacts can be found at <http://drought.unl.edu/Planning/Impacts.aspx>

meteorological quantity associated with dry conditions, namely, precipitation. In addition, we also consider conditions in near surface air temperature, which can affect surface drying through increased evaporative demand in warmer air, although the latter can also result from soil drying associated with meteorological drought itself (e.g. Mueller and Seneviratne 2012; Sheffield et al. 2012; Yin et al. 2014). We begin with an overview of the interannual variability of both precipitation and temperature.

Figure 1 shows the land regions where SST anomalies are expected to influence annual mean precipitation and two meter (2 m) air temperature, based on the five Atmospheric General Circulation Models (AGCMs: 12 ensemble members for each) forced with observed SST over the period 1979-2011. (See Appendix A for descriptions of the models and simulations). The base maps show the fraction of the total interannual variance that is forced by SST. Focusing on precipitation (Fig. 1a), we see that the ratios outside the tropics (poleward of  $\pm 30^\circ$  latitude) are generally quite small ( $< 0.2$ )<sup>3</sup>; outside the tropics, much of the interannual variability is unforced by SST and is therefore likely to be unpredictable from SST forcing at interannual time scales. This is for instance the case in northern Eurasia, central Europe as well as central and eastern Canada. We note nonetheless that (agricultural and hydrological) drought predictability in these regions may be arising from year-to-year memory in soil moisture and/or snow pack, or possibly interannual changes in radiative forcing, aspects that we do not consider in the present review. The largest fractions of interannual precipitation variability explained by SST in the midlatitudes occur over the U.S. southern Great Plains,

---

<sup>3</sup> We note that values of the ratio greater than 0.06 are statistically significant at the 1% level based on a F-test following Zwiers et al. (2000).



189 southwest Asia, parts of Australia and South America. Values exceed 0.3 primarily in  
190 tropical land areas, including northwest South America, Indonesia, Central America,  
191 southeast Asia, southwestern India, and eastern Africa. The fractions for 2m temperature  
192 (Fig. 1b) are generally considerably larger than those for precipitation. Some regions in  
193 the extratropics show values exceeding 0.4 (e.g., southern US Great Plains and Mexico).  
194 Nevertheless, the largest values are again confined to the tropical regions of Africa,  
195 southern Asia, Indonesia and much of the northern half of South America, with values  
196 sometimes exceeding 0.7.

197 El Niño/Southern Oscillation (ENSO) is a key player in the development of precipitation  
198 deficits in many regions of the world (e.g., Ropelewski and Halpert 1987). Figure 1, in  
199 addition to showing the fraction of interannual variance forced by SST, shows how SST  
200 is correlated with precipitation (Fig. 1a) and T2m (Fig. 1b) within selected regions (small  
201 inserts)<sup>4</sup>; the patterns show a clear link to ENSO and to SST in general. We will refer to  
202 these maps as we review the results from the individual contributions to this special  
203 collection.

204 A number of regions of the world have suffered multi-year drought (e.g., beyond the  
205 ENSO time scale), and one may wonder whether such droughts result from naturally  
206 occurring decadal modes of variability (e.g., the Atlantic Multi-decadal Oscillation or

---

<sup>4</sup> We emphasize that these are meant to be summary results. As we shall see in section 3, there are in some cases considerable variations in SST connections within a box and between seasons. For example, the western portions of East Africa tend to have a JJAS rainfall maximum, and El Nino is tied to drought. Further east (eastern Ethiopia, Kenya, Somalia) the rainy season is bimodal with drought associated with La Nina (and its influence on Indian Ocean SSTs) during boreal fall. The ENSO signal reverses sign between East and Southern Africa as well, with 15° south frequently considered the northern limit of the southern African region. Some of these seasonal and regional differences are discussed in section 3.

207 AMO and Pacific Decadal Oscillation or PDO), from decadal changes in the relationships  
208 between interannual modes of variability (e.g., ENSO and Atlantic Niños; Losada et al.  
209 2012), from global warming (Mohino et al. 2011a), or from no mechanism at all, i.e.,  
210 from a simple random sequence of dry years generated from internal atmosphere  
211 variability. In Figure 2 we provide a global depiction of the changes that have occurred  
212 over the last three decades in the tails of the probability distributions of 2m temperature  
213 and precipitation based on the same set of AGCM runs used to produce the results in  
214 Figure 1. Here we show how the probability of exceeding (or falling short of, in the case  
215 of precipitation) a particular critical value  $x_c$  has changed between the first and last  
216 fifteen years of the record. Because we are focusing on extreme years,  $x_c$  is chosen to be  
217 the 2.5% value based on all 33 years – i.e., the value that would be exceeded (or fallen  
218 short of) on average only 2.5% of the time. The last three decades, we note, are  
219 characterized by both global warming and shifts in the AMO and PDO (Figure 3), so  
220 anthropogenic forcing and natural variations may both contribute significantly to  
221 observed regional changes between these two periods.

222

223 In regard to precipitation (Fig. 2a), the models indicate that much of the United States has  
224 experienced an increase in the probability of extreme dry years during the last three  
225 decades (particularly the central Plains). Here the shift is 1 to 1.5 times the  
226 climatological probability of 2.5%. The shift is clear in the probability density functions  
227 (pdfs) provided in the insert. As we shall see next, this shift reflects forcing by SST with  
228 a strong decadal component and does not necessarily reflect a long-term trend. In fact, if

a longer time period is considered, the United States (especially the central part of the country) has generally experienced wetter conditions compared to the 1950s (e.g., Wang et al. 2009; Seneviratne et al. 2012; Hartmann et al. 2013; Greve et al. 2014). As Wang et al. 2009 showed however, even for this longer time period the precipitation “trend” is still dominated by SST forcing with decadal time scales.

Parts of Indochina and southeastern China also see an increase in the probability of extreme dry years. In contrast, northeastern South America shows a substantial decrease in the probability of dry years over the last three decades, though with little change in the probability of extreme wet years (see insert). The tropical west coast of Africa, the Sahel, and northeastern Russia also show a reduction in the probability of extreme dry years. The pdf characterizing precipitation in northeastern Africa shows no shift in the peak, so that the changes in the pdf occur primarily in the tails. In general, for the Northern Hemisphere during the last three decades, the high latitudes show a tendency for a reduction in the probability of dry years, whereas the middle latitudes (including parts of Europe, southern Asia and the U.S.) show a tendency for an increase in the probability of dry years. In the Southern Hemisphere, the probability of extreme dry years is increased in parts of southern Africa, Australia and southern and western South America and mostly decreased in tropical regions.

Relative to precipitation, the results for 2m temperature (Figure 2b) are more homogenous, with almost all regions of the world showing an increase in the probability of very warm years over the last three decades (see also Hartmann et al. 2013). Regions where the increase in probability exceeds twice the climatological probability of 2.5%

251 include the south central U.S., Mexico, northwestern South America, eastern Canada,  
252 parts of Europe, southern Asia, Japan, tropical and northern Africa, Indonesia, and  
253 southern Australia. Only northeastern South America and western Canada show  
254 substantial regions with little increase (and even some scattered regions of decrease) in  
255 the probability of warm years. The inserts show that these changes largely result from a  
256 shift in the mean rather than from a change in the shape of the pdfs for the analyzed  
257 regions.

258 Figure 3 compares the simulated and observed mean changes between the two 15 year  
259 periods. The model results show warming everywhere except over northwestern North  
260 America and northeastern South America, with the strongest warming occurring in the  
261 Northern Hemisphere. The model results are generally consistent with the observed  
262 temperature changes, although they are smoother due to being an average over 60  
263 ensemble members. There are also strong similarities between the simulations and  
264 observations in the precipitation differences, with both difference maps showing  
265 decreases over the U.S. and increases over northern South America, northern Australia,  
266 northern Eurasia, and central Africa. Some differences in the estimated precipitation  
267 changes, however, do appear, including over central South America (observed decreases  
268 not found in the simulations), India, and southeast Asia. The extent to which these reflect  
269 model deficiencies or sampling differences associated with unforced internal atmospheric  
270 noise is unclear. Overall, the changes are consistent with the changes in the pdfs  
271 discussed earlier. They appear to reflect, in part, a response to SST changes linked to the

PDO, the AMO, and a warming trend (Fig. 3, see also Schubert et al. 2009), as well as possible direct impacts on the atmosphere from increasing GHGs<sup>5</sup>.

### **3. Causes of Meteorological Drought by Region**

We now provide a more in-depth discussion of meteorological drought for specific regions. While much of the discussion is condensed from the individual contributions to this special collection, we also present relevant results from the aforementioned SST-forced AGCM simulations, as well as results from other key studies where necessary to address issues not covered by the individual contributions.

We begin by providing in Figure 4 a brief assessment of the ability of the models to produce the observed annual cycle of precipitation in each of the selected regions (see the boxes in Figures 5-8 for the definitions of the regions). This is also meant to facilitate the following discussion about the links to SST, by giving the reader an assessment of the timing of the wet and dry seasons in each region. Overall, the models do a reasonable job in reproducing the observed annual cycle, though the peak rainfall tends to be underestimated especially in the tropical land regions (N. South America/Central America, Southern Eurasia, Indonesia)<sup>6</sup>. It is noteworthy that the Central/South American region (Figure 4a) shows some evidence of the well-known mid summer drought found over Mexico and Central America (e.g., Magaña et al. 1999) – something that is also reproduced in the model results. We also note that the spatial averaging tends to hide any regional differences. This is especially true for the east African region which

---

<sup>5</sup> All the AGCMs (except for CCM3) were forced with observed GHGs (Appendix A).

<sup>6</sup> We note that including the ocean points when computing the area averages of the precipitation in these regions produces much closer agreement between the observations and model results (not shown), indicating the underestimation of the precipitation is confined to the land areas.

shows a rather flat annual cycle (Figure 4b), despite having local rainfall regimes that include unimodal (JJA and DJF maxima) and bimodal (MAM, OND maxima) annual cycles (see section 3.3).

### *3.1 North America*

The occurrence of precipitation deficits over North America on annual time scales is predominantly associated with SST variability in the tropical Pacific (e.g., Seager et al. 2005), with some contribution from SST variability in the Atlantic (e.g., Schubert et al. 2009). Figure 1a shows that precipitation deficits are largely tied to La Nina conditions, with the largest impacts in the southern Great Plains and Northern Mexico. La Nina conditions also lead to warming across the Southern Plains and much of the southeast, whereas El Nino conditions are associated with warming over Alaska and northwestern Canada (Fig. 1b).

These results are consistent with the in-depth assessment of the causes of North American drought carried out by Seager and Hoerling (2014). Using a subset of the climate models underlying Figure 1, Seager and Hoerling (2014) find that SST forcing of annual mean precipitation variability accounts for up to 40% of the total variance in northeastern Mexico<sup>7</sup>, the southern Great Plains and the Gulf Coast states but less than 10% in central and eastern Canada. They further find that, in addition to the tropical Pacific, tropical North Atlantic SST contributes to the forcing of annual mean

---

<sup>7</sup> Mexico will be discussed further in the following section.

precipitation and soil moisture in southwestern North America and the southern Great Plains.

Seager and Hoerling (2014) find that SST forcing was indeed responsible for multiyear droughts in the 1950s and at the turn of the 21<sup>st</sup> Century. Attribution to SST patterns, however, is not always straightforward. Wang et al. (2014) highlight how the responses over North America to SSTs in different ocean basins can reinforce each other or cancel out, complicating the analysis of SST impacts. Atmospheric internal variability also muddies the signal; internal atmospheric variability can contribute significantly to extreme droughts, especially on shorter (monthly) time scales (Seager et al 2014a; Hoerling et al. 2014; Wang et al. 2014). For example, the most extreme phase of the Texas drought in 2011 was largely unforced by SST, and the central Plains drought of 2012 showed almost no contribution from SST forcing.

Figure 5 (left panels) shows, for the United States and northern Mexico, the seasonality of the link between precipitation and SST, as determined from the model simulations. The most striking aspect of this seasonality is the strong ENSO connection for all seasons except JJA, though the strong connection in MAM is not supported by the observations (see Fig. B5). Summertime precipitation is negatively correlated with tropical Atlantic SST, a result consistent with Kushnir et al. (2010) and Wang et al. (2008), who showed that a larger Atlantic Warm Pool leads to a suppressed Great Plains Low Level jet and associated reduced Central U.S. precipitation. On the other hand, summertime precipitation is positively (though weakly) correlated with SST along the west coast of

North America extending into the central tropical Pacific, with a structure reminiscent of the PDO. The link to the Indian Ocean also has substantial seasonality, with positive correlations during DJF and MAM and negative correlations extending westward from the warm pool into the eastern Indian Ocean during SON.

Seager and Hoerling (2014) show that, during the early 21st Century, natural decadal variations in tropical Pacific and North Atlantic SSTs have contributed to a dry regime for the U.S. (see also Fig. 3). Since the mid 1990s both the PDO and the AMO have gone through striking decadal transitions (Figure 3) to a cold tropical Pacific-warm North Atlantic that is “ideal” for North American drought (Schubert et al. 2009). Figure 2 indicates that in the southern plains region, the drier regime is associated with a substantial increase in the probability of extreme dry years. In addition, Seager and Hoerling (2014) note that long-term changes caused by increasing trace gas concentrations are now contributing to a modest signal of soil moisture depletion, mainly over the American Southwest, thereby prolonging the duration and severity of naturally occurring droughts.<sup>8</sup>

Understanding the extent to which precipitation and air temperature variability is determined by SST forcing (potentially providing predictability) and internal atmospheric variability (providing no predictability on seasonal and longer time scales) is an important research challenge (e.g., Wang et al. 2014). Recently the 2011-14 California drought has been linked to a localized warm SST anomaly in the western tropical Pacific

---

<sup>8</sup> Water pumping is another source of drying in the SW US.



(Seager et al. 2014c; Hartmann 2015), which raises the important issue of forcing of drought over North America by Pacific SST anomaly patterns other than ENSO. The contribution of soil moisture to the variability is also still poorly understood, as reflected by the substantial differences in the strength of land-atmosphere feedback and soil moisture memory simulated by current climate models (e.g., Koster et al. 2004; Seneviratne et al. 2006). Also poorly understood is the nature and predictability of the unforced component (e.g., internal atmospheric variability associated with Rossby waves and other atmospheric teleconnections, especially during the summer).

Regarding changes under enhanced greenhouse gas concentrations and global warming, the additional forcing of increasing radiation could lead to enhanced evapotranspiration during drought events. Climate projections for the end of the 21<sup>st</sup> century suggest a robust increase of soil moisture drying in the southern United States and Mexico, while signals for accumulated precipitation deficits are less robust across climate models (Orlowsky and Seneviratne 2013). However, historical records do not yet suggest a detectable signal in North America, either in precipitation or precipitation-evapotranspiration (Hartmann et al. 2013; Greve et al. 2014). How the SST impacts may change in a warming world is largely unknown.

### *3.2 Latin America*

Figure 1 shows that SST impacts on temperature and precipitation are strong over northern South America; these signals are largely associated with ENSO and tropical Atlantic variability (e.g., Mechoso and Lyons 1988; Saravanan and Chang 2000; Giannini

et al. 2004). Via this connection, this region may see substantial improvements in seasonal prediction skill as climate models improve (e.g., Folland et al. 2001; Goddard et al. 2003). In Central America, as in northern South America, precipitation is correlated negatively with tropical Pacific SST and positively with tropical Atlantic SST. Indeed, the extent to which the Atlantic signals are independent of ENSO is still not fully quantified (e.g., Chang et al. 2003). Extreme droughts in northeast Brazil have been linked to very strong El Niño events (McCarthy et al. 2001). Conversely, western Amazon droughts depend on tropical North Atlantic SST anomalies more than on ENSO (Marengo et al 2008). Further analysis demonstrated that the North Tropical Atlantic influence is largest during dry season droughts in the southern Amazon, but ENSO still has a stronger influence during the wet season for the entire basin (Yoon and Zeng 2009).

Figure 1 also shows that relatively strong signals for precipitation over South America extend south along the west coast which shows enhanced precipitation associated with La Niña conditions. Relatively high temperature signals along the west coast extending southward into northern Chile are associated with positive correlations with El Niño. The east coast over southern Brazil and Uruguay, including northern and central Argentina (much of the La Plata basin), has reduced precipitation associated with La Niña conditions (Diaz et al. 1997; Fig. 1a). According to McCarthy et al. (2001), during La Niña events Chile and central-western Argentina, exhibit negative anomalies of rainfall and snowfall leading to reduced summer streamflow.

Figure 5 illustrates the seasonality of the link to SST over northern South America and

405 Central America (middle column) and over central-southern South America (right  
406 column). For the former region, the aforementioned link to the El Niño cycle is weakest  
407 in March-April-May (MAM), and the link to the tropical Atlantic is strongest during  
408 June-July-August (JJA) and September-October-November (SON). On the other hand,  
409 Cazes-Boezio et al. (2003) show that the ENSO impact on precipitation in Uruguay  
410 occurs primarily during austral spring (October–December), but is almost absent during  
411 peak summer (January –February), followed by weak impacts during March-July. This is  
412 not inconsistent with our much larger central-southern South American region (and with  
413 somewhat different definitions of the seasons), which is characterized by reduced  
414 (enhanced) precipitation in association with La Niña (El Niño) conditions for all seasons  
415 except December-January-February (DJF), when correlations with SST are negligible.  
416

417 While droughts in southeastern South America exhibit a strong dependence on La Niña  
418 (cold Pacific), a warm North Tropical Atlantic can help define the shape and intensity of  
419 the drought episodes (Seager et al. 2010; Mo and Berbery 2011). Notably, the effect of  
420 land surface-atmosphere interactions, in the form of soil moisture-precipitation coupling,  
421 is essential in the development of drought in southern South America (Xue et al. 2006;  
422 Wang et al. 2007; Ma et al. 2010; Sörensson and Menéndez 2011). Barreiro and Diaz  
423 (2011) noted that improved seasonal forecasts over South America require the proper  
424 representation of the teleconnection processes and regional land–atmosphere interactions  
425 need to be adequately resolved. Müller et al. (2014) showed that during the severe 2008  
426 drought in southern South America, a realistic representation of land surface biophysical  
427 properties leads to a better depiction of surface-atmosphere processes that consequently

reduces model biases and eventually contributes to improved prediction skill of droughts.

McCarthy et al. (2001) note that in Central America, topography influences the ENSO impacts; during El Niño years, the Pacific side is characterized by reduced precipitation, while some parts of the Caribbean side have above normal rain. They also note that over Colombia, El Niño events are associated with reductions in precipitation, streamflow, and soil moisture, whereas La Niña is associated with heavier precipitation and floods (Poveda and Mesa, 1997), especially during December-January. El Niño also tends to bring large positive precipitation anomalies to the eastern part of the Andes, Ecuador, and northern Peru.

Future climate scenarios produced by regional downscaling suggest a precipitation decrease over the tropical region of South America, with an increase over the subtropical areas (Sanchez et al. 2015). In relation to extremes, climate change scenarios for South America suggest an increase of dry spells, with more frequent warm nights (Marengo et al. 2009).

### 3.3 *East Africa*

Lyon (2014) provides a review of the regional and large-scale SST and atmospheric circulation patterns associated with meteorological drought in East Africa on seasonal and longer time scales. Analysis of drought in the region is complicated by local rainfall regimes that generally consist of unimodal (JJA and DJF maxima) and bimodal (MAM, OND maxima) annual cycles. On seasonal-to-interannual time scales, ENSO is the largest source of seasonal rainfall variations, but depending on season and location, it has opposite effects: La Niña is frequently associated with drought during the OND “short rains” in the central and eastern areas of the Greater Horn (this is not well captured by most of the models – see Figures 6 and B1), whereas El Niño is linked to deficient rainfall during boreal summer in locations further west having a unimodal annual cycle (consistent with Figure 6). Particularly for the short rains, the Indian Ocean plays a critical role in mediating the impact of ENSO, with the development of a west-east Indian Ocean SST anomaly dipole pattern (IOD) being closely associated with rainfall variations (see also Figure 6). ENSO, however, is associated with at most roughly 25% of interannual variations in East African rainfall (consistent with Figure 1).

In observations, interannual variations in MAM “long rains” (Funk et al. 2008; Lyon and DeWitt 2012) in East Africa do not show statistically significant correlation with SSTs in any ocean basin (generally consistent with Figure 6, though the models do show positive correlations with SST in the western Indian Ocean). At longer time scales, AMIP-style model runs do tend to capture the decline in the East African long rains associated with the shift in Pacific SSTs towards the cool phase of the PDO in 1998-99 (Lyon 2014,

Yang et al. 2014, see also Figures 3 and 9). The models may thus respond more to decadal, rather than interannual, variations in SSTs. Liebmann et al. (2014) suggest this result may be tied to the relative magnitudes of multidecadal SST fluctuations relative to interannual variability.

On longer timescales, there is growing concern over an observed increase in the frequency of drought, primarily during the MAM long rains. This increase has had dire impacts across the Greater Horn, with the most recent and severe drought in 2010-11 helping to trigger a humanitarian crisis and contributing to the fatalities of tens of thousands of people. The increase in drought frequency has raised concerns about the possible role of anthropogenic climate change. Paradoxically, the consensus of climate model projections is for the region to become wetter during the current century in response to anthropogenic greenhouse gas forcing (IPCC 2007). Lyon et al. (2013), Lyon (2014) and Yang et al. (2014) provide evidence that the recent rainfall decline is substantially driven by natural, multi-decadal variability, a result consistent with our model simulations (Figure 9). Consistent with Lyon and DeWitt (2012), subsequent studies by Hoell and Funk (2013, 2014) suggest that long-term anthropogenic warming of the western Pacific may further enhance the equatorial SST gradient associated with the cold phase of the PDO and thus also enhance drying in East Africa during MAM. As to whether East Africa will become wetter or drier as a result of anthropogenic forcing, Yang et al. (2014) caution that most coupled climate models do not properly capture either the observed annual cycle of rainfall in East Africa or the observed relationship between seasonal rainfall variations and SSTs in different basins (particularly the Indian

Ocean), calling into question the reliability of climate projections in East Africa. Lyon (2014) concludes that the hydroclimatic response of East Africa to anthropogenic climate change remains an open question and that more research is needed to better understand the physical processes associated with rainfall variability of the region across multiple timescales.

### *3.4 West Africa and Sahel*

Rodríguez-Fonseca et al. (2015) focus on rainfall variability across multiple timescales in West Africa and the Sahel. They conclude that SST variations are largely responsible for rainfall variability in the region. Land surface processes and aerosols including those from volcanic eruptions modulate the SST influence.

The left column in Fig. 6 indicates a strong seasonality in the correlation between West African (including western Sahel) rainfall and SST in the simulation by the five models with prescribed SST corresponding to the observed over the period 1979-2011. During the rainy season (JJA), increased precipitation over West Africa is associated with colder SST in the eastern tropical Pacific and northern Indian Ocean, and with warmer SST in the tropical Atlantic/Gulf of Guinea. During the dry season (DJF), when climatological precipitation is small, increased precipitation is associated with warmer SST in the tropical Atlantic/Gulf of Guinea, as well as in the tropical North Atlantic and central tropical Pacific. Correlations are weaker and less organized in the Pacific during MAM, and little connection with SST is apparent during SON.

Other experiments using AGCMs with prescribed SSTs in individual ocean basins have provided additional insight. During the wet season, warm equatorial SST anomalies corresponding to a warm Atlantic Niño (Rodríguez Fonseca et al. 2009) are associated with precipitation increases over the Gulf of Guinea and weaker decreases over the Sahel. The impact of a Pacific warm event varies during the season. In the early part of the season (May-June) warming of the equatorial Pacific reduces rainfall over the Gulf of Guinea and enhances it over the Sahel. In the late part of the wet season (July-August) warming of the equatorial Pacific reduces rainfall over the Sahel. In the seasonal mean, the negative effects of the Pacific Niños in the late season prevail over the positive ones in the early period as shown in Fig. 6.

Rodríguez-Fonseca et al. (2015) discuss a unique aspect of the West African rainfall variability at interannual time scales: its links with the variability of tropical SSTs have shown non-stationary features (see also Rodríguez-Fonseca et al. 2011 and Mohimo et al. 2011b). Pacific cold events and Atlantic warm events tend to appear simultaneously after the 1970's (Rodríguez-Fonseca et al. 2009). AGCM experiments demonstrate that, during this period, the impacts of simultaneous SST anomalies in the Indo-Pacific and Atlantic on Sahel rainfall tend to cancel each other such that the north-south dipole in rainfall anomalies over West Africa expected from Atlantic SST anomalies only does not appear in the observations (Losada et al. 2012).

Analysis of observational data and model results has provided clues on the mechanisms at work in the connections described above. Anomalous warming of the southern tropical



Atlantic enhances ascent over the Gulf of Guinea and descent over the Sahel. A warming in the Pacific and Indian Oceans generates equatorial Rossby waves that contribute to subsidence over the Sahel and thus to reduce regional precipitation. Also, Mediterranean warm events are linked to increased moisture flux convergence over the Sahel.

Decadal SST variability and global warming are also relevant to Sahelian drought. In recent decades the Sahel has been recovering from a devastating drought in the 1970s and 80s. It has been suggested that a special combination of three different modes of SST variability (the global warming trend, the positive phase of the Inter-decadal Pacific Oscillation, and the negative phase of the Atlantic Meridional Oscillation, or AMO) led to this drought (Mohino et al., 2011a). Vegetation dynamics has been contributing to regional climate persistence (e.g., Zeng et al. 1999). The recovery from the drought appears to be driven by SST also, as a similar feature is obtained in SST-forced model simulations. Regarding global warming, Rodríguez-Fonseca et al. (2015) note that, while rainfall projections have a large spread, models do show a tendency for slightly wetter conditions over the central Sahel and drier conditions over the west. The onset of the rainy season is projected to be delayed, especially over West Africa, while more abundant precipitation is expected during the late rainy season.

Rodríguez-Fonseca et al. (2015) caution that more research is needed to further support these model-based findings on the variability of Sahel rainfall. While most models capture, for example, the link with Mediterranean SSTs, some important teleconnections are still not well reproduced (e.g., those linked to equatorial Atlantic SSTs and Pacific ENSO; Rowell 2013). Also, coupled atmosphere-ocean general circulation models have great difficulties in correctly

reproducing the seasonal cycle and variability of the tropical Atlantic SST (including the Atlantic Equatorial mode; Richter 2015) and Pacific (e.g. Mechoso et al. 1995).

### *3.5 The Middle East and Southwest Asia*

Barlow et al. (2015) provide a comprehensive review of current understanding of drought in the Middle East and Southwest Asia – a region that is water-stressed, societally vulnerable, and prone to severe droughts. They note that this understanding is still at an early stage, though it appears that large-scale climate variability, particular La Niña in association with a warm western Pacific, contributes to region-wide drought, including the two most severe droughts of the last fifty years (1999-2001 and 2007-2008). Barlow et al. (2015) provide a schematic for those two years indicating how La Niña-related SSTs and a warm western Pacific led to wave responses that affected vertical motion, moisture flux and storm tracks in the region. They note that the North Atlantic Oscillation (NAO), the AMO, and the Atlantic SST tripole pattern also influence the region, though the strength of the teleconnections varies considerably within the region, and the temporal stability of the relationships is somewhat uncertain.

Figure 1 (top panel) highlights the role of ENSO (and perhaps the PDO) in influencing drought in Southwest Asia on annual time scales. This result shows some model dependence but appears to be consistent with observations (Figure B2). Figure 7 (left column) shows that there is a strong seasonality to the precipitation-SST connection in this region, with the strongest correlations in MAM (La Niña, together with a cool tropical Indian Ocean and cool tropical North Atlantic, is apparently conducive to

drought conditions then) and similar, though somewhat weaker (especially in the tropical Indian Ocean) correlations in DJF. These two seasons comprise the wet season for most of the region, associated with synoptic precipitation (Barlow et al. 2015). Warm season precipitation is important in Pakistan and the southern coast of the Arabian Peninsula, associated with the Indian monsoon and the ITCZ. During JJA the link to SST changes sign in the tropical Pacific, so that reduced precipitation is linked to warm tropical Pacific SSTs together with cold SSTs in the tropical Atlantic. SON shows the beginnings of the cold-season link to ENSO, with a coherent ENSO pattern extending from the western Mediterranean into Southwest Asia (Mariotti 2007).

Barlow et al. (2015) note that in the high mountains of the region, snowmelt provides predictability for peak river flows and potentially for vegetation; vegetation in the region is closely linked to precipitation and may also play a feedback role. The drying of the eastern Mediterranean is a robust feature of future projections, as are temperature increases across the region.

### *3.6 East Asia*

Zhang and Zhou (2015) review drought over East Asia with a primary focus on China. They point out that due to the seasonal variation of monsoonal circulation, drought mostly occurs over North China and Southwest China in spring, with the highest drought frequency and maximum duration occurring during that season. In early July, drought tends to occur in the Yangtze River and Huaihe River valleys of China and also Korea and Japan due to the influence of the Northwestern Pacific subtropical High.

606

607 The interannual variability of East Asian summer (EASM) precipitation is in part  
608 associated with the Pacific-Japan teleconnection pattern, which features a meridional tri-  
609 polar pattern during decaying El Nino summers, with excessive precipitation in central  
610 eastern China along the Yangtze River valley (27.5°N-32.5°N,102°E-120°E) but drier or  
611 even drought conditions in southern (20°N-27.5°N,102°E-120°E) and northern (32.5°N-  
612 45°N,102°E-120°E) China, or vice versa (Huang and Li et al. 2007). This is associated  
613 with an anomalous anticyclone over the western North Pacific forced by the SST  
614 anomalies there and over the Indian Ocean during decaying El Niño summers (Li et al.  
615 2008; Xie et al. 2009; Wu et al. 2009, 2010). The Silk Road teleconnection, a pattern  
616 forced by Indian monsoon heating and characterized by the propagation of a stationary  
617 Rossby wave along the Asian jet in the upper troposphere, also affects East Asia, mainly  
618 North China precipitation (Wu et al. 2003; Ding et al. 2011).

619

620 Drought trends over China since about 1950 are characterized by a zonal dipole pattern,  
621 with an increasing trend over the central part of North China and a decreasing tendency  
622 over Northwest China. The drying and warming trend over North China is associated  
623 with an inter-decadal weakening of the East Asian summer monsoon circulation, which  
624 has been mainly linked to the 1970s phase transition of the PDO from negative to  
625 positive values (Zhou et al. 2009a, 2013). While the weakening of the monsoon  
626 circulation is well reproduced by AMIP-type simulations (Li et al. 2010), the associated  
627 anomalous precipitation change found in observations is poorly reproduced over East  
628 Asia. This is likely in part due to the bias' that exist in simulating the climatological

precipitation in this region, resulting from the inability of current relatively coarse global models to resolve the complex terrain over Asia (Zhou et al. 2008a,b, Li et al. 2015).

While CMIP5 experiments indicate that aerosols act to weaken monsoon circulation, the simulated change is much weaker than observed (Song et al. 2014). A 50-70 year variation in the PDO index appears to be imprinted in century-scale variations of drought in North China (Qian and Zhou 2014).

Up to now, most efforts have focused on exploring the prediction skill of East Asian monsoon precipitation, with few examining drought predictability. Previous studies show that climate models have limited skill in simulating and predicting the precipitation in terms of both climatological mean state and interannual variations (Chen et al. 2010; Zhou et al. 2009b). In contrast, the variability of East Asian monsoon circulation is well captured (Zhou et al. 2009c; Song and Zhou 2014a). A successful reproduction of the interannual EASM pattern depends highly on the Indian Ocean–western Pacific anticyclone teleconnection (Kim et al. 2012; Song and Zhou 2014a,b). Finally, Zhang and Zhou (2015) note that in climate change projections, most climate models simulate an increasing drought frequency and intensity over East Asia, mainly in southeastern Asia, though the models do differ regarding drought patterns and severity.

Figure 7 (rightmost column) shows that the link between precipitation over eastern China, Korea, and Japan with SST varies seasonally, with the strongest ties in DJF and MAM – reduced precipitation in the region is tied to La Nina and cold Indian Ocean SST.

652 During JJA the correlation to SST is overall weak. During SON the correlations with  
653 precipitation are negative in the western North Pacific and positive in the northern Indian  
654 Ocean and the eastern tropical Pacific.

655

656 These results are consistent with those of Yang and Lau (2004), who found that MAM  
657 precipitation in southeastern China is linked to ENSO – reduced precipitation occurs in  
658 years with an abnormally cold central and eastern tropical Pacific and Indian Ocean.

659 Yang and Lau (2004) found that on average in southern China (south of the Yangtze  
660 River), MAM and JJA precipitation each account for about 35% of the annual total, with  
661 JJA presenting a more complicated picture (see above). They also found that in years  
662 with abnormally warm SSTs over the warm pool and northern Indian Ocean and  
663 abnormally cold SSTs over the western North Pacific, precipitation over central eastern  
664 China tends to be anomalously high (see also Wu et al. [2009, 2010]). They further found  
665 a tendency for a weakened East Asian monsoon circulation and a delayed monsoon onset  
666 in years for which SSTs in the central and eastern tropical Pacific are abnormally warm,  
667 resulting in reduced late summer precipitation over northern China.

668

669 The above linkages are reflected to some extent in the model results of Figure 7 (right  
670 panels), though without any evidence of a strong link to SST in the warm pool. Figure 7  
671 in fact features the typical SST anomaly patterns that dominate the East Asian climate  
672 during the mature phase of El Niño (boreal winter) extending into the decaying-year  
673 summer (Wu et al. 2009). This type of interannual monsoon-SST relationship is well  
674 captured by the AMIP simulations of CMIP3 and CMIP5 models (Song and Zhou 2014a).

675

676 A comparison of the observed and model generated changes in Figure 3 indicates that the  
677 reduced precipitation over southeast China over the last three decades is linked to SST.  
678 Figure 9 shows these long-term changes have occurred primarily during spring and fall,  
679 though DJF does show an enhanced probability of extreme dryness (Fig. 10). Figure 11  
680 shows that the warming of the last three decades is associated with an enhanced  
681 probability of extreme warm seasons especially during JJA over northwest China, Korea  
682 and Japan, and over most of East Asia during SON.

683

### 684 *3.7 India*

685 Kanikicharla et al. (2015), in their comprehensive review of monsoon droughts over  
686 India, note that Indian drought is indeed synonymous with monsoon failure and that a  
687 number of historical droughts there have led to severe famines and great human and  
688 economic losses. They use a century-long time series of Indian summer monsoon rainfall  
689 (ISMR) to capture the characteristic spatio-temporal features of deficit monsoons and  
690 their possible driving mechanisms. They particularly discuss the low-frequency  
691 modulation of ISMR and associated drought area extent in India with respect to global  
692 climate phenomena, and they employ a large suite of AMIP-type model simulations to  
693 assess the predictability of Indian drought.

694

695 Some key findings from that paper are:

696

-Monsoon failures are linked to preceding winter and spring snow accumulation over the Himalayas and larger regions of Eurasia and to the occurrence of warm ENSO events in the Pacific (with the latter link being much stronger).

-The leading EOF of Indian monsoon rainfall has a very conspicuous resemblance to the rainfall anomaly pattern associated with major droughts, and that EOF's time series correlates well with an ENSO-like SST pattern in the Pacific.

- The low frequency behavior of monsoon rainfall and drought area index goes hand in hand with the opposite sign of the NIÑO34 index (which captures the ENSO and AMO), though with a large difference in their evolution in recent decades. This indicates that the behavior of the Indian monsoon in recent decades cannot be fully explained by known global teleconnections and that other factors (e.g., Indian Ocean variability, aerosols) could be influencing its variability on interannual and decadal time scales (e.g., Ramanathan et al. 2005; Lau et al. 2006; Gautam 2009).

- AGCM simulations forced with global and regional SSTs are able to reproduce the low-frequency variability well, and runs with observed SSTs in the Pacific but with climatological SSTs elsewhere generally produce the sign of many droughts in the past century. The simulated rainfall deficits, however, are much smaller than observed



- More concerted efforts with climate models are needed to anticipate the severity and geographical extent of droughts. Global warming is probably altering known teleconnections, complicating our ability to predict Indian drought.

These findings emphasize the challenges faced in predicting drought over India and surrounding regions within a changing climate. Figure 1 (top panel) emphasizes the weak link of annual mean precipitation over southern Asia to global (and in particular ENSO) SST in recent decades, though it also shows (bottom panel) that temperature variations in the broader south Asian monsoon region do have strong ties to global SST. Figure 7 (middle column of panels) highlights the strong seasonality in the simulated link between precipitation and SST in recent decades, with only MAM showing a substantial link to ENSO: El Niño (La Niña) is associated with reduced (enhanced) precipitation. (This link is consistent with observations and robust across the models [Figure B3].) ENSO may provide important preconditioning of the land (e.g., soil moisture, snow) during the pre-monsoon months, so that the role played by SST in monsoon droughts, while important, may be indirect. We note that such effects may be missed by contemporaneous correlations, as in the present analyses.

### *3.8 Australia and the Maritime Continent*

As reviewed in Cai et al. (2014), the influence of climate variability and change on Australia is complex and varies both regionally and seasonally. In particular, they indicate how the continent is impacted by the Indian Ocean Dipole (IOD), the Southern

Annular Mode (SAM), and El Niño-Southern Oscillation (ENSO), and the poleward edge of the Southern Hemisphere Hadley cell.

The corresponding correlation map in Figure 1a shows aspects of a negative IOD, La Niña and a negative PDO. Consistent with the discussion in Cai et al. (2014), the key SST forcing regions driving Australian precipitation appear to be the tropical Pacific just west of the dateline and areas in the eastern Indian Ocean just to the north and west of Australia. This is highlighted in Figure 8 (left panels), with spring (SON) and summer (DJF) showing the strongest relationship between Australian precipitation and remote SST. In summer, the pan-Australian rainfall is dominated by contributions from northern and eastern Australia; as such, dry conditions are associated with both a warm central tropical Pacific (with weaker correlations extending eastward across the Pacific), i.e., an El Niño, and a warm Indian Ocean SST (basin-wide warming that usually accompanies an El Niño). In spring, the contributions to the pan-Australian rainfall come about equally from northern and southern Australia, and ENSO and the IOD have the highest coherence to rainfall in those regions. As such, spring appears to have the strongest (spatially most coherent) link to ENSO, with dry conditions linked to El Niño and a cool anomaly in the eastern Indian Ocean. (This result is robust across models and observations [Figure B4].) In contrast, during fall (MAM) and particularly during winter (JJA), the pan-Australian precipitation comes mostly from southern Australia. The deficit during fall (MAM) shows the greatest link with cold SST anomalies to the northwest, while the deficit during winter (JJA) is linked to cold SST anomalies in the eastern Indian Ocean associated with

the development phase of a positive IOD. These cold SST anomalies are unfavorable for weather systems that typically deliver rain-producing moisture over southern Australia.

Recently, Australia experienced one of its most severe droughts ever recorded: the “Millennium Drought”, which lasted about 10 years (2000-2009). Cai et al. (2014) showed that the associated precipitation anomalies had substantial seasonal variation, with austral summer (DJFM) showing positive precipitation anomalies in northwest Australia and with some of the largest deficits over other parts of Australia occurring during late fall and winter. Figure 3 shows that the annual mean differences (1998-2011 minus 1979-1993) largely reflect the summer precipitation increases in northwest Australia during that drought. The relevant seasonality is well captured by the models (Figure 9), which show the northern Australian precipitation surfeits in the recent 15-year period during DJFM and the deficits associated with the Millennium Drought during AMJJ (cf. Figure 3 of Cai et al. [2014]). Much of Australia in fact experienced an increased risk of dry winters (JJA) over the last fifteen years (Fig. 10).

Cai et al. (2014) show that the precipitation deficits over southwest Western Australia partly result from a long-term upward trend in the Southern Annual Mode (SAM); this trend accounts for half of the winter rainfall reduction there. For southeast Australia, CMIP5 model simulations indicate only weak trends in the pertinent climate modes, apparently underestimating the observed poleward expansion of the subtropical dry-zone and associated impacts. They conclude that “*although climate models generally suggest that Australia’s Millennium drought was mostly due to multi-decadal variability, some*

late-twentieth-century changes in climate modes that influence regional rainfall are partially attributable to anthropogenic greenhouse warming.”

The Maritime Continent is strongly affected by ENSO during JJA and SON (Figure 8, right panels); El Niño conditions lead to reduced precipitation. JJA also exhibits positive correlations with tropical and North Atlantic SST. In contrast, DJF and MAM precipitation show little connection with ENSO, and the overall correlations with SST are weak – weak negative correlations with SST in the central tropical Pacific, and, for MAM, weak positive correlations with local SST. Chang et al. (2003), however, point out that the low correlations between Indonesian rainfall and ENSO during the Northern Hemisphere winter monsoon period are partly due to the spatial averaging of the rainfall in two regions with opposite characteristics.

### *3.9 Europe and the Mediterranean*

Here we review the primary modes of variability that affect European and Mediterranean climate on subseasonal to interannual and longer time scales, with a focus on their impacts on precipitation and/or surface temperature fields. We shall see that northern and central European meteorological drought drivers and impacts are often different or even opposite to those for southern Europe and the Mediterranean region<sup>9</sup>. In addition, we will discuss reported trends in meteorological drought in Europe and projected drought changes in Europe with increasing greenhouse gases.

---

<sup>9</sup> In this section, the term Mediterranean is used to indicate areas surrounding the sea from southern Europe, northern Africa and the Middle East; the term Europe indicates northern and central Europe.

807 Correlations of the North Atlantic Oscillation (NAO) and the Atlantic Multidecadal  
808 Oscillation (AMO) with drought occurrence in Europe have been documented, and the  
809 effects of other modes of variability including ENSO have been postulated (see below).  
810 Nonetheless, these relationships do not seem to be associated with high interannual  
811 predictability of meteorological drought in central and northern Europe (Dutra et al.  
812 2014). Overall SST anomalies, which may be associated with large-scale modes of  
813 variability, explain only a small fraction of annual mean variability in precipitation (less  
814 than 10%) and temperature (less than 20%) over Europe (see Fig. 1). Hence the  
815 predictability associated with large-scale modes of variability that have been linked to  
816 drought occurrence in Europe is still unclear from the existing literature. In addition, it  
817 has been highlighted that the circulation patterns and weather types related to the most  
818 severe droughts in Europe often vary across seasons and regions (Stahl 2001, see also  
819 below; Fleig et al. 2011).

820

821 Hurrell (1995) showed that during high NAO index winters (such as those that occurred  
822 in 1983, 1989 and 1990), the axis of maximum moisture transport shifts to a more  
823 southwest-to-northeast orientation across the Atlantic and extends much farther north and  
824 east onto northern Europe and Scandinavia, accompanied by a reduction in moisture  
825 transport over southern Europe, the Mediterranean and northern Africa. As a result,  
826 northern Europe is mild and wet during the positive phase of the NAO and cold and dry  
827 during the negative phase, whereas the reverse is true for southern Europe and most of  
828 the Mediterranean but with the Levant also being wet (dry) during a positive (negative)  
829 NAO (e.g. Xoplaki et al. 2004). In recent decades the NAO index has shown a return

toward more negative values, though with marked increase in year-to-year winter variability (Hanna et al. 2015).

The summer NAO (sNAO) has a more northerly location and smaller spatial scale than its winter counterpart (Folland et al. 2009). Nevertheless, the sNAO has a strong influence on northern European rainfall, temperature, and cloudiness through changes in the position of the North Atlantic storm track, thus playing an important role in generating summer climate extremes, including flooding, drought, and heat waves in northwestern Europe. A positive sNAO also results in wetter conditions in the central/eastern part of the Mediterranean.

Folland et al. (2009) further suggest that on interdecadal time scales, sNAO variations are partly related to the Atlantic Multidecadal Oscillation (AMO). While the exact link between the two is still unclear, Sutton and Dong (2012) show that, during the 1990s, European climate shifted towards a pattern characterized by anomalously wet summers in northern Europe and hot, dry, summers in southern Europe, with related shifts in spring and autumn, and they point to recent evidence suggesting that the warming was largely caused by an acceleration of the Atlantic Meridional Overturning Circulation (AMOC) and associated northward ocean heat transport in response to the persistent positive phase of the winter NAO in the 1980s and early 1990s (Robson et al. 2012). However, uncertainties still exist regarding the processes underpinning AMO variability, for example the role of anthropogenic aerosols (Booth et al. 2012). Mariotti and dell'Aquila (2012) show that decadal variability associated with the NAO, the sNAO, and the AMO

significantly contribute to decadal climate anomalies over the Mediterranean region. The positive phase of the AMO is associated with warmer than usual decades especially in summer, whereas the AMO has no influence on Mediterranean winter temperatures. Land-atmosphere feedbacks also play role in shaping observed decadal variability, enhancing the large-scale influences. Della Marta et al. (2007) found a relationship between western Mediterranean heat waves and the AMO. On decadal timescales, the AMO and NAO explain over 60% of observed area-averaged summer temperature and winter precipitation variability, respectively (Mariotti and Dell'Aquila, 2012).

The Mediterranean is one of the main regions worldwide displaying a robust drying trend in both precipitation and the land water balance since the 1950s (Sheffield et al. 2012; Hartmann et al. 2013; Greve et al. 2014), a signal consistent with climate change projections (see below). Nonetheless, a possible attribution of these historical trends to increased greenhouse gas concentrations has not been provided so far, and it is possible that decadal variability associated with large-scale modes of variability could have played a role. Hoerling et al. (2012) note that for the land area surrounding the Mediterranean Sea, 10 of the 12 driest winters since 1902 occurred in just the last 20 years, and they propose that the drying over the last century can be understood as a response to a uniform global ocean warming and to modest changes in the oceans' zonal and meridional SST gradients, with warming in the Indian Ocean producing an enhanced drying signal attributable to an atmospheric circulation response resembling the positive phase of the NAO. Kelley et al. (2011), in a combined observational and modeling analysis, argue that while the upward NAO-trend over recent decades can explain much of the

concurrent Mediterranean region drying, it cannot explain drying in the Levant which they instead argued was consistent with drying in response to rising greenhouse gases.

With respect to climate change projections, the Mediterranean shows one of the most robust responses (across models) to greenhouse gas increases over the 21<sup>st</sup> Century (Giorgi and Lionello 2008; Seneviratne et al. 2012; Orłowsky and Seneviratne 2013). Projected changes, which reinforce trends already observed during the 20<sup>th</sup> Century, include both a reduction in precipitation and an increase in evapotranspiration (due to increased incoming radiation and higher air temperature), with associated soil moisture reductions (e.g. Mariotti et al. 2008; Orłowsky and Seneviratne 2012, 2013; Seager et al. 2014b; Mariotti et al. 2015).

Uncertainties remain in our understanding of the NAO's link to SST changes (Bretherton and Battisti 2000) and of its link to global warming (Gillett et al. 2003), its connection to the Arctic Oscillation (AO, Ambaum et al. 2001), its link to the stratosphere (Scaife et al. 2005), and its possible modulation by ENSO (Bönnimann 2007) and other modes of variability such as the Scandinavian (SCA) and the East Atlantic (EA) patterns (e.g., Comas-Bru and McDermott 2014). In fact, the SCA, the EA, and the East Atlantic/western Russia (EAWR) patterns (Barnston and Livezey 1987) have also been suggested to contribute substantially to European climate variability (e.g. Bueh and Nakamura 2007; Iglesias et al. 2014; Ionita 2014).



A number of studies have produced objectively-defined drought catalogues for Europe (Lloyd-Hughes et al. 2010; Spinoni et al. 2015). Parry et al. (2010) also produced summaries of the major large-scale European droughts of the last 50 years based on the catalogue from Lloyd-Hughes et al. (2010). As summarized in Stahl (2001), major droughts over the period 1960-1990 occurred during 1962-64, 1972-76, 1983, 1989-1990. They note that “*Most of the severe summer droughts across Europe were associated with high pressure systems across central Europe. Generally, drought associated synoptic meteorology is characterised by high MSLP, but the circulation pattern types vary not only with the season but also for all individual discussed events.*”

Figures 2b and 11 show that there has been a pronounced shift in the probability of extremely warm years over the last three decades over most of Europe, with the shift equal to more than 1.5 times the climatological probability of 2.5% over many regions. This shift, most pronounced during fall, appears to be associated with a shift in the mean temperature over the recent decades, which is likely attributable in part to enhanced greenhouse gas forcing (Bindoff et al. 2013). Figures 2a and 10 show that changes in precipitation over the last three decades are generally small, though there is a general tendency for a greater probability of extremely dry years throughout central and southern Europe. This appears to hold for each season as well. This result is also robust when investigating longer-term trends since the 1950s, either for precipitation or precipitation minus evapotranspiration (Seneviratne et al. 2012a; Hartmann et al. 2013; Greve et al. 2014).

While the present review and special issue focus on meteorological (i.e. precipitation-based) drought and its relation to SSTs as driver, we note the following additional points regarding drought drivers in Europe:

- In general, agricultural (soil moisture) and hydrological (streamflow) drought events in Europe are caused by a prolonged deficit in precipitation (Tallaksen et al. 2015, Stagge et al. 2015). However, in Central Europe, evapotranspiration is an important driver for soil moisture droughts, in some cases to the same degree as precipitation (e.g. Seneviratne et al. 2012b, Teuling et al. 2013). In addition, in cold climates, temperature anomalies also play a role in the development of hydrological drought (Van Loon and Van Lanen 2012).

- Prior storage deficits in the form of soil moisture, snow and groundwater are important for the occurrence and development of soil moisture and streamflow droughts (van Loon and van Lanen 2012, Orth and Seneviratne 2013, Staudinger et al. 2014, Tallaksen et al. 2015). They thus provide some essential sources of drought predictability, in particular given the low SST control on interannual precipitation variability in Europe (Fig. 1).

### *3.10 Northern Eurasia*

Figure 1 shows that across the vast expanse of Northern Eurasia, neither precipitation nor temperature is strongly affected by SST on interannual time scales. Schubert et al. (2014), in examining both heat waves and drought over northern Eurasia, highlighted the central role of anticyclones in the region; these act to warm and dry the atmosphere and

land surface over many important agricultural regions from European Russia to Kazakhstan and beyond. They discuss how the development of anticyclones is linked to different air masses, especially the intrusion of Arctic air masses that occasionally combine with subtropical air (e.g., associated with the Azores high in eastern Europe and western Russia) to produce especially severe drought and heat wave events. Schubert et al. (2014) found that some of the most severe summer heat waves are linked to distinct Rossby wave trains spanning the continent that, while producing severe heat in one location, cause a juxtaposition of wet and cool conditions in regions thousands of miles to the east or west – a phenomenon noted more than one hundred years ago in early descriptions of Northern Eurasian heat waves.

Given the lack of a strong SST connection, the predictability of the most severe events in Northern Eurasia is limited to the time scales of the internally forced Rossby waves (typically less than one month), though some aspects of heat waves appear to be predictable for several months: the surface temperature anomalies at the center of the heat wave associated with soil moisture anomalies that persist through the summer. Schubert et al. (2014), using an AGCM experiment in which soil moisture feedbacks were disabled, showed that temperature variability is strongly tied to soil moisture variability particularly in the southern parts of Northern Eurasia extending from southern Europe eastward across the Caucasus, Kazakhstan, Mongolia, and northern China. They note that longer-term droughts (lasting multiple years) do occur but are largely confined to the southern parts of Northern Eurasia, where there appears to be a weak link to SST and an important control from soil moisture.

967  
 968 Schubert et al. (2014) further showed that the observed warming over northern Eurasia in  
 969 the last three decades is part of a large-scale warming pattern with local maxima over  
 970 European Russia and over Mongolia/eastern Siberia (see also Fig. 3). Precipitation  
 971 changes consist of deficits across Eurasia covering parts of northeastern Europe,  
 972 European Russia, Kazakhstan, southeastern Siberia, Mongolia, and northern China and  
 973 increases across Siberia poleward of about 60°N. Comparisons of these changes with  
 974 Figure 3, however, indicate some sensitivity of the computed changes to the years chosen  
 975 for averaging. Model simulations carried out with idealized versions of the observed SST  
 976 anomalies indicate that the changes over the last three decades are consistent with a  
 977 global-scale response to PDO-like and AMO-like SST patterns, intensified by a global  
 978 warming SST trend.  
 979  
 980 Figure 2 indicates that any changes in probability of heat waves are likely a consequence  
 981 of an overall warming trend that affects much of Eurasia (though more strongly in the  
 982 southern regions). In particular, the increase in the probability of extreme heat largely  
 983 results from an overall shift in the pdf of temperature (a change in the mean) rather than  
 984 from any change in the shape of that pdf. Schubert et al. (2014) point to studies  
 985 indicating an enhanced probability of heat waves across northern Eurasia by the second  
 986 half of the 21<sup>st</sup> Century. Existing studies and analyses of climate change projections of  
 987 the Coupled Model Intercomparison Project (CMIP), however, show less certainty  
 988 regarding future drought (e.g. Seneviratne et al. 2012; Orlowsky and Seneviratne 2013),  
 989 reflecting the greater uncertainty of precipitation and soil moisture projections.

990

#### 991 **4. Concluding Remarks**

992 The results presented here, and in the regionally-focused articles that make up this special  
993 collection, illustrate that considerable progress has been made in our understanding of the  
994 occurrence and predictability of meteorological drought in different parts of the world.

995 The importance of large-scale teleconnections, whether they are linked to ENSO or other  
996 SST variability or whether they consist of large-scale atmospheric circulation anomalies  
997 that are unforced by SST (internal to the atmosphere), is now well established. As such,  
998 in addressing the causes and predictability of meteorological drought for any particular  
999 region of the world, we have to address the following questions: 1) what are the large-  
1000 scale drivers (if any) of precipitation deficits relevant to that region, and 2) what are the  
1001 unique climatological features of that region that act to enhance or suppress the large  
1002 scale precipitation tendencies.

1003

1004 Although the individual articles have in many cases provided answers to both of these  
1005 questions and we have attempted to summarize them in Section 3, this article goes further  
1006 by providing a more global perspective on these two questions in the context of the  
1007 “consensus” view provided by the simulations with current state-of-the-art AGCMs  
1008 forced by observed SST.

1009

1010 In particular, we have provided our current best estimate of where SST forcing provides  
1011 some control on annual precipitation and temperature variability. This is critically  
1012 important to the drought prediction problem, since the regions where SST do play a

substantial role in driving precipitation (and temperature) variability, are also the regions where we can expect to have some degree of predictability on seasonal and longer time scales. We have also underscored the importance of ENSO (tropical Pacific SST) in providing that potential predictability in many parts of the world (including the Americas, eastern and southwest Asia, Australia and the Maritime Continent) though not exclusively so, with other Ocean basins also playing a role in some regions of the world, either individually or in concert with ENSO. These include the Indian Ocean (the Indian Ocean dipole affecting Australia, the Indian Ocean basin mode affecting East Asia), the Atlantic Ocean (affecting northern South America, parts of the southern and eastern US, and the Sahel), and the Mediterranean Sea (affecting southern Europe and northern Africa), though the extent to which some of these SST patterns are independent of ENSO is still not fully resolved.

A number of regions dominated by monsoonal climates have droughts that are intimately tied to failures in the development of monsoonal rains. The GDIS papers highlight the substantial progress made in identifying the sources of these failures. From a global perspective, ENSO significantly affects much of the Asian-Australian monsoon system. On decadal time scales, the apparent weakening of the global land monsoon since the 1950s has been linked to the Inter-decadal Pacific Oscillation as well as to a warming trend over the central-eastern Pacific and the western tropical Indian Ocean. Much remains to be understood, however, about the observed trends in monsoonal precipitation.

1035 Northern Eurasia, central Europe, and central and eastern Canada stand out as regions  
1036 with little SST-forced impacts on precipitation on interannual time scales. This has  
1037 important implications for the predictability and the time scales of droughts in these  
1038 regions. In Northern Eurasia, for example, droughts and heat waves are predominantly  
1039 linked to the development of anticyclones, and although extreme, they rarely last longer  
1040 than one season. In central Europe a number of different atmospheric teleconnections  
1041 that are unforced (or only weakly forced) by SST do appear to play a role, though these  
1042 are relatively short lived and have little predictability on seasonal and longer time scales:  
1043 here evapotranspiration is an important driver for soil moisture droughts, and  
1044 predictability on longer time scales is tied to soil moisture memory and feedbacks.

1045

1046 Although the annual mean results provide a broad picture of the role of SST, our results  
1047 also highlight the strong seasonality in the link to the SST that occurs in most regions of  
1048 the world. As such, the timing and duration of drought has as much to do with the  
1049 seasonality of the link to SST, as it has to do with the time of year that local  
1050 climatological (land, circulation) conditions make a region most prone to drought. East  
1051 Africa is an example of a particularly challenging region for which to model and  
1052 understand drought, due to heterogeneous local rainfall regimes that include unimodal  
1053 and bimodal annual cycles combined with strong seasonality in the response to ENSO.

1054

1055 We have also addressed longer-term (decadal) meteorological drought and the link to  
1056 SST. In particular, we present a remarkable example of the ability of current climate  
1057 models (when forced with observed SST) to reproduce the long-term changes in

precipitation and surface temperature that have occurred over the last three decades. The model results show that the shifts to drought or pluvial conditions (and warming) have a global coherence driven by long-term SST changes (a combination of the PDO, AMO, and a long term trend). Our analysis of the most extreme seasonal and annual mean precipitation deficits shows that the associated changes in the tails of the probability density functions (PDFs) in most regions of the world reflect the overall shifts in the mean rather than changes in the shape of the PDFs (though this may not be true for northeastern South America – a region exhibiting a substantial decrease in the probability of extremely dry years over the last three decades though with little change in the probability of extreme wet years). The success of the models in reproducing the observed changes provides the basis for further research to dissect the causes of these changes and address their potential predictability.

One consequence of the decadal and longer-term variations is that a number of regions exhibit substantial non-stationarity in the relationships between SST and precipitation on interannual time scales (examples where this is particularly evident include west Africa and India), complicating our ability to understand these relationships, and take advantage of them for prediction. Global warming, while not a focus of this paper, is clearly an important issue when addressing longer-term changes in drought. In fact, as discussed in a number of the contributed papers, some regions of the world appear to already be seeing the impacts of warming on drought (e.g., the southwestern U.S., the Mediterranean region and central Europe), though much work needs to be done to better understand the relative contributions of decadal SST variability and long term SST trends to drought.



1081

1082 Finally, we must emphasize that current climate models, including the AGCMs used  
1083 here, are far from perfect. A key factor emphasized in many of the contributing papers  
1084 and further highlighted here is the challenge of reproducing some of the complex local  
1085 precipitation regimes (including the annual cycle) that must be simulated correctly in  
1086 order to properly simulate the impact of large scale forcing on regional drought. The  
1087 relatively coarse resolution of current climate models hinders that, and so we can at best  
1088 obtain a spatially averaged picture or in some cases even an incorrect assessment of the  
1089 impacts. Examples where this is especially critical include East Africa and East Asia –  
1090 regions that are characterized by complex terrain and highly heterogeneous regional  
1091 precipitation regimes. We should note that this situation will likely improve in the  
1092 coming years as it becomes feasible to apply ultra-high resolution (sub 10 km) global  
1093 models to climate problems. However, not all problems concerning the simulation of  
1094 important teleconnections can be blamed on insufficient resolution. For example,  
1095 deficiencies in the simulation of the atmospheric response to equatorial Atlantic SSTs and  
1096 the link to west African drought are likely tied to deficiencies in the simulation of the  
1097 climatological mean state. Furthermore, considerable work still needs to be done to  
1098 improve our coupled atmosphere-ocean general circulation models that still have, for  
1099 example, great difficulties in correctly reproducing the seasonal cycle and variability of  
1100 tropical Atlantic SST.

1101

1102 How do we move forward? Drought is an immensely complex problem that must be  
1103 attacked on many fronts by researchers from around the world, with well-considered

links to users who may benefit from the research. GDIS is an ongoing activity that supports this cause. GDIS will continue to encourage researchers and users around the world to work together to improve systematically our understanding of, prediction of, and adaptation to drought, e.g., by facilitating the development of improved models and long term consistent drought-specific observations, and providing global access to data portals that summarize our ever-evolving knowledge on the subject.

*Acknowledgements:* The various contributions to this paper were made possible by the support of the host organizations of the co-authors, as noted in the acknowledgments of the contributing Global Drought Information System (GDIS) special collection papers. The GDIS effort is sponsored and supported by the World Climate Research Program (WCRP: CLIVAR and GEWEX) and various partner organizations including the US National Oceanic and Atmospheric Administration (NOAA), the US National Aeronautics and Space Administration (NASA), the US National Integrated Drought Information System (NIDIS), the World Meteorological Organization (WMO), the US CLIVAR program, the Group on Earth Observations (GEO), the European Commission Joint Research Centre (JRC), the National Science Foundation (NSF), and the European Space Agency (ESA)-European Space Research Institute (ESRIN). Support for the overall development of this synthesis paper was provided by NASA's Modeling, Analysis and Prediction Program. The GLDAS-2 data used in this study were acquired as part of the mission of NASA's Earth Science Division and archived and distributed by the Goddard Earth Sciences (GES) Data and Information Services Center (DISC).

1127

1128

## Appendix A: Description of the Models and Experiments

Many of the results presented in this paper are based on Atmospheric Model Intercomparison Project (AMIP)-style simulations produced with 5 different Atmospheric General Circulation Models (AGCMs). The models used are GEOS-5, CCM3, CAM4, GFS, and ECHAM5. *The years 1979-2011 were subsetted from 12 of each model's ensemble members, providing a dataset of 60 33-yr simulations.* The following provides a brief description of the models and the experiments.

The NASA Goddard Earth Observing System Model, Version 5 (GEOS-5) Atmospheric General Circulation model (AGCM) is described by Rienecker et al. (2008), and an overview of the model's performance is provided by Molod et al. (2012). For these experiments, the model was run with 72 hybrid-sigma vertical levels extending to 0.01hPa and with a 1° horizontal resolution on a latitude/longitude grid. The simulations consist of twelve ensemble members, each forced with observed monthly SST, sea ice, and time-varying greenhouse gases for the period 1871-present. See Schubert et al. (2014) for further details.

A 16-member ensemble covering the period January 1856 to April 2014 was produced with the NCAR Community Climate Model 3 (CCM3, Kiehl et al. (1998)). The model was run at T42 resolution with 18 vertical levels. Sea ice was held at climatological values, and SST forcing in the years of interest here combined the Kaplan et al. (1998) SST dataset in the tropical Pacific Ocean (20°N to 20° S) with the Hadley Centre dataset (Rayner et al. 2003) outside of the tropical Pacific.

1152

1153 A 20-member ensemble covering the period January 1979 to April 2014 was produced  
1154 with the NCAR Community Atmosphere Model 4 (CAM4), forced by SST and sea ice  
1155 from the Hurrell et al. (2008) data set and with time varying GHGs from the RCP6.0  
1156 scenario after 2005. The resolution used was  $0.94^{\circ} \times 1.25^{\circ}$ , with 26 vertical levels.

1157

1158 The NOAA Earth System Research Laboratory produced a 50-member ensemble  
1159 spanning January 1979 to April 2014 using the NCEP Global Forecast System (GFS, the  
1160 atmosphere component of the Coupled Forecast System) version 2 model denoted here as  
1161 (ESRL GFSv2). The model was run at T126 resolution with 64 vertical levels and was  
1162 forced by observed SST and sea ice from the Hurrell et al. (2008) data. CO<sub>2</sub> varied with  
1163 time, but other GHGs were held fixed.

1164

1165 A 20-member ensemble spanning January 1979 through April 2014 was produced with  
1166 the ECHAM5 model (Roeckner et al. 2003) forced by the Hurrell et al. (2008) SST and  
1167 sea ice data, as recommended for use in CMIP5 simulations. These simulations used  
1168 time-varying GHGs, following the RCP6.0 scenario after 2005, and they used a T159  
1169 resolution, with 31 vertical levels.

1170

1171

## **Appendix B: Selected Individual Model Results**

Here we present a few comparisons of the results for individual models and observations, highlighting regions where it is especially important to assess the model-dependence of the results (see main text). While the comparison with observations provides a rough idea of consistency with nature, it must be kept in mind that the observations represent a single realization of nature and thus should differ from the ensemble means of the model runs, which specifically isolate the impact of SST and other forcings – our focus here. A careful assessment of the veracity of the models, which is beyond the scope of this paper, would in principle involve determining if a correlation produced for the observations lies within the spread produced by the given model’s individual ensemble members.

*(Figure B1 here)*

*(Figure B2 here)*

*(Figure B3 here)*

*(Figure B4 here)*

*(Figure B5 here)*

## References

- Ambaum, M. H. P., B. J. Hoskins, and D. B. Stephenson, 2001: Arctic Oscillation or North Atlantic Oscillation? *J. Climate*, **14**, 3495–3507.  
doi: [http://dx.doi.org/10.1175/1520-0442\(2001\)014<3495:AOONAO>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(2001)014<3495:AOONAO>2.0.CO;2)
- Barlow, M., B. Zaitchik, S. Paz, E. Black, J. Evans, A. Hoell, 2015: A Review of Drought in the Middle East and Southwest Asia. *J. Climate*. Under review.
- Barnston A.G., Livezey R.E., 1987: Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Mon. Wea. Rev.*, **115**, 1083–1126.
- Barreiro, M., N. Diaz, 2011: Land-atmosphere coupling in El Niño influence over South America. *Atmos. Sci. Lett.*, **12**, 351-355, doi: 10.1002/asl.348.
- Bindoff, N.L., P.A. Stott, K.M. AchutaRao, M.R. Allen, N. Gillett, D. Gutzler, K. Hansingo, G. Hegerl, Y. Hu, S. Jain, I.I. Mokhov, J. Overland, J. Perlwitz, R. Sebbari and X. Zhang, 2013: Detection and Attribution of Climate Change: from Global to Regional. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

1212 Booth, B. B. B., N. J. Dunstone, P. R. Halloran, T. Andrews, and N. Bellouin, 2012:  
 1213 Aerosols implicated as a prime driver of twentieth-century North Atlantic climate  
 1214 variability. *Nature*, 484, 228–232.  
 1215

1216 Bretherton, C. S. and D. S. Battisti, 2000: An interpretation of the results from  
 1217 atmospheric general circulation models forced by the time history of the observed sea  
 1218 surface temperature distribution. *Geophys. Res. Lett.*, 27, 767-770.  
 1219

1220 Brönnimann, S., 2007: Impact of El Niño-Southern Oscillation on European climate.  
 1221 *Rev. Geophys.* 45, RG3003, doi: 10.1029/2006RG000199.  
 1222

1223 Bueh C. and H. Nakamura, 2007: Scandinavian pattern and its climatic impact. *Q. J. R.*  
 1224 *Meteorol. Soc.* 133 2117–31.  
 1225

1226 Cai, W., A. Purich, T. Cowan, P. van Rensch, E. Weller, 2014: Did Climate Change-  
 1227 Induced Rainfall Trends Contribute to the Australian Millennium Drought? *J. Climate*,  
 1228 27, 3145 – 3168.  
 1229

1230 Cazes-Boezio, G., A. W. Robertson, and C. R. Mechoso, 2003: Seasonal dependence of  
 1231 ENSO teleconnections over South America and relationships with precipitation in  
 1232 Uruguay. *J. Climate*, 16, 1159–1176.  
 1233



1234 Chang, P., R. Saravanan, and L. Ji, 2003: Tropical Atlantic seasonal predictability: The  
1235 roles of El Niño remote influence and thermodynamic air–sea feedback. *Geophys. Res.*  
1236 *Lett.*,30,1501, doi:10.1029/2002GL016119.

1237

1238 Chen, H., T. Zhou, R. B. Neale, X. Wu, G. Zhang, 2010: Performance of the New NCAR  
1239 CAM3.5 in East Asian Summer Monsoon Simulations: Sensitivity to Modifications of  
1240 the Convection Scheme. *J. Climate*, 23, 3657-3675.

1241

1242 Comas-Bru, L. and F. McDermott, 2014: Impacts of the EA and SCA patterns on the  
1243 European twentieth century NAO–winter climate relationship. *Q. J. R. Meteorol. Soc.*,  
1244 140, 354–363, January 2014B.

1245

1246 Cook, B.I., T.R. Ault, and J.E. Smerdon, 2015: Unprecedented 21st-century drought risk  
1247 in the American Southwest and Central Plains. *Sci. Adv.*, **1**, no. 1, e1400082,  
1248 doi:10.1126/sciadv.1400082.

1249

1250 Dai, A., K. E. Trenberth, and T. Qian, 2004: A global data set of Palmer Drought  
1251 Severity Index for 1870-2002: Relationship with soil moisture and effects of surface  
1252 warming. *J. Hydrometeorology*, 5, 1117-1130.

1253

1254 Della-Marta, P. M., J. Luterbacher, H. von Weissenfluh, E. Xoplaki, M. Brunet, H.  
1255 Wanner, 2007: Summer heat waves over western Europe 1880-2003, their relationship to  
1256 large-scale forcings and predictability. *Clim. Dyn.*, 29, 251-275. DOI 10.1007/s00382-

1257 007-0233-1.

1258

1259 Diaz, A. F., C. D. Studzinski and C. R. Mechoso, 1997: Relationships between

1260 precipitation anomalies in Uruguay and southern Brazil and sea surface temperature in

1261 the Pacific and Atlantic oceans. *J. Climate*, 11, 251-271.

1262

1263 Ding, Q., and B. Wang, J. M. Wallace, and G. Branstator, 2011: Tropical–extratropical

1264 teleconnections in boreal summer: Observed interannual variability. *J. Climate*, 24,

1265 1878–1896, doi: 10.1175/2011JCLI3621.1.

1266

1267 Dutra et al., 2014: Global meteorological drought – Part 2: Seasonal forecasts. *Hydrol.*

1268 *Earth Syst. Sci.*, 18, 2669-2678.

1269

1270 Fleig, A.K., L.M. Tallaksen, H. Hisdal, and D.M. Hannah, 2011: Regional hydrological

1271 drought in north-western Europe: Linking a new Regional Drought Area Index with

1272 weather types. *Hydrological Processes*, 25, 1163–1179.

1273

1274 Folland, C. K., Colman, A. W., Rowell, D. P. and Davey, M. K. 2001. Predictability of

1275 northeast Brazil rainfall and real-time forecast skill, 1987–98. *J. Climate*, 14, 1937–1958.

1276

1277 Folland, Chris K., Jeff Knight, Hans W. Linderholm, David Fereday, Sarah Ineson, and

1278 James W. Hurrell, 2009: The Summer North Atlantic Oscillation: Past, Present, and

1279 Future. *J. Climate*, **22**, 1082–1103. doi: <http://dx.doi.org/10.1175/2008JCLI2459.1>

1280

1281 Funk, C., M. D. Dettinger, J. C. Michaelsen, J. P. Verdin, M. E. Brown, M. Barlow, and  
1282 A. Hoell, 2008: Warming of the Indian Ocean threatens eastern and southern African  
1283 food security but could be mitigated by agricultural development. *Proc. Natl. Acad. Sci.*  
1284 USA, **105**, 11081–11086, doi:10.1073/pnas.0708196105.

1285

1286 Gautam, R., [N. C. Hsu](#), [W. Lau](#), and M. Kafatos (2009), Aerosol and rainfall variability  
1287 over the Indian monsoon region: distributions, trends and coupling, *Ann. Geophys.*, **27**,  
1288 3691-3703.

1289

1290 Giannini, A., Saravanan, R. and Chang, P. 2004. The preconditioning role of tropical  
1291 Atlantic variability in the development of the ENSO teleconnection: implications for the  
1292 prediction of Nordeste rainfall. *Clim. Dyn.* **22**, DOI: 10.1007/s00382-004-0420-2, 839–  
1293 855.

1294

1295 Gillett, N. P., F. W. Zwiers, A. J. Weaver, and P. A. Stott, 2003: Detection of human  
1296 influence on sea-level pressure, *Nature*, **422**, 292–294.

1297

1298 Giorgi, F., and P. Lionello, 2008: Climate change projections for the Mediterranean  
1299 region. *Global and Planetary Change*, **63**, 90-104.

1300

1301 Goddard, L., Barnston, A. G. and Mason, S. J. 2003. Evaluation of the IRI's 'net  
1302 assessment' seasonal climate forecasts: 1997–2001. *Bull. Amer. Meteor. Soc.* **84**, 1761–

1303 1781.

1304

1305 Greve, P., B. Orlowsky, B. Mueller, J. Sheffield, M. Reichstein, and S.I. Seneviratne,

1306 2014: Global assessment of trends in wetting and drying over land. *Nature Geoscience*, 7,

1307 716-721, doi: 10.1038/NGEO2247.

1308

1309 Hanna, E., T.E. Cropper, P.D. Jones, A.A. Scaife, R. Allan, 2015: Recent seasonal

1310 asymmetric changes in the NAO (a marked summer decline and increased winter

1311 variability) and associated changes in the AO and Greenland Blocking Index. *Int. J.*

1312 *Climatol.*, 35, 2540-2554.

1313

1314 Hartmann, D.L., A.M.G. Klein Tank, M. Rusticucci, L.V. Alexander, S. Br. nnimann, Y.

1315 Charabi, F.J. Dentener, E.J. Dlugokencky, D.R. Easterling, A. Kaplan, B.J. Soden, P.W.

1316 Thorne, M. Wild and P.M. Zhai, 2013: Observations: Atmosphere and Surface. In:

1317 Climate Change 2013: The Physical Science Basis. Contribution of Working Group

1318 I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change

1319 [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y.

1320 Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United

1321 Kingdom and New York, NY, USA.

1322

1323 Hartmann, D., 2015: Pacific sea surface temperature and the winter of 2014. *Geophys.*

1324 *Res. Let.* Doi:10.1002/2015GL063083

1325

1326 Hoell, A., and C. Funk, 2013: The ENSO-related West Pacific sea surface temperature  
 1327 gradient. *J. Climate*, 26, 9545-9562, doi:10.1175/JCLI-D-12-00344.1.  
 1328  
 1329 \_\_\_\_\_, and \_\_\_\_\_, 2014: Indo-Pacific sea surface temperature influences on failed  
 1330 consecutive rainy seasons over eastern Africa. *Climate Dyn.*, 43, 1645-1660,  
 1331 doi:10.1007/s00382-013-1991-6.  
 1332  
 1333 Hoerling, M. P., and A. Kumar, 2003: The perfect ocean for drought. *Science*, 299, 691-  
 1334 694.  
 1335  
 1336 Hoerling, Martin, Jon Eischeid, Judith Perlwitz, Xiaowei Quan, Tao Zhang, and Philip  
 1337 Pegion, 2012: On the Increased Frequency of Mediterranean Drought. *J. Climate*, 25,  
 1338 2146–2161. doi: <http://dx.doi.org/10.1175/JCLI-D-11-00296.1>  
 1339  
 1340 Hoerling, M., J. Eischeid, A. Kumar, R. Leung, A. Mariotti, K. Mo, S. Schubert, and R.  
 1341 Seager, 2014: Causes and Predictability of the 2012 Great Plains Drought. *Bull. Amer.*  
 1342 *Meteor. Soc.*, 95, 269–282. doi: <http://dx.doi.org/10.1175/BAMS-D-13-00055.1>  
 1343  
 1344 Hurrell, J. W., 1995: Decadal trends in the North Atlantic Oscillation: regional  
 1345 temperatures and precipitation. *Science*, 269, 676-679.  
 1346

1347 Hurrell, James W., James J. Hack, Dennis Shea, Julie M. Caron, and James Rosinski,  
 1348 2008: A New Sea Surface Temperature and Sea Ice Boundary Dataset for the Community  
 1349 Atmosphere Model. *J. Climate*, 21, 5145–5153.  
 1350 doi: <http://dx.doi.org/10.1175/2008JCLI2292.1>  
 1351  
 1352 Iglesias I., Lorenzo M. N., Taboada J. J., 2014: Seasonal Predictability of the East  
 1353 Atlantic Pattern from Sea Surface Temperatures. *PLoS ONE* 9(1): e86439. doi:10.1371/  
 1354 journal.pone.0086439  
 1355  
 1356 Ionita, M., 2014: The Impact of the East Atlantic/Western Russia Pattern on the  
 1357 Hydroclimatology of Europe from Mid-Winter to Late Spring. *Climate*, 2(4), 296-309.  
 1358 doi:10.3390/cli2040296.  
 1359  
 1360 IPCC, 2001: *Climate Change 2001: Impacts, Adaptation and Vulnerability*, Contribution  
 1361 of Working Group II to the Third Assessment Report of the Intergovernmental Panel on  
 1362 Climate Change. [McCarthy, J. J.; Canziani, O. F.; Leary, N. A.; Dokken, D. J.; and  
 1363 White, K. S., ed.], Cambridge University Press, ISBN 0-521-80768-9 (pb: 0-521-01500-  
 1364 6).  
 1365  
 1366 IPCC, 2007: *Climate Change 2007: The Physical Science Basis*. Contribution of Working  
 1367 Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate  
 1368 Change [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M.

1369 Tignor and H.L. Miller (eds.)). Cambridge University Press, Cambridge, United  
 1370 Kingdom and New York, NY, USA, 996 pp.  
 1371  
 1372 IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working  
 1373 Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate  
 1374 Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A.  
 1375 Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press,  
 1376 Cambridge, United Kingdom and New York, NY, USA, 1535 pp,  
 1377 doi:10.1017/CBO9781107415324.  
 1378  
 1379 Kanikicharla, K. K., A. Kulkarni, S. Pai and S. Nigam, 2015: Monsoon droughts in India,  
 1380 J. Climate, under review.  
 1381  
 1382 Kaplan, A., M. A. Cane, Y. Kushnir, A. C. Clement, M. B. Blumenthal, and B.  
 1383 Rajagopalan, 1998: Analyses of global sea surface temperature 1856– 1991. J. Geophys.  
 1384 Res., 103, 18,567– 18,589.  
 1385  
 1386 Kelley, C., M. Ting, R. Seager and Y. Kushnir, 2011: The relative contributions of  
 1387 radiative forcing and internal climate variability to the late 20th Century winter drying of  
 1388 the Mediterranean region. Clim. Dyn., 38, 2001-2015.  
 1389

1390 Kiehl J. T., Hack J. J., Bonan G. B., Boville B. A., Williamson D. L., Rasch P. J., 1998:  
 1391 The National Center for Atmospheric Research Community Climate Model: CCM3. J.  
 1392 Climate, 11, 113-1149.  
 1393  
 1394 Kim H. M., P. J. Webster, J. A. Curry and V. E. Toma, 2012: Asian summer monsoon  
 1395 prediction in ECMWF System 4 and NCEP CFSv2 retrospective seasonal forecasts.  
 1396 Climate Dyn., 39(12), 2975-2991.  
 1397  
 1398 Koster, R. D. and Coauthors, 2004: Regions of strong coupling between soil moisture and  
 1399 precipitation. Science, 305, 1138–1140.  
 1400  
 1401 Kushnir, Y., R. Seager, M. Ting, N. Naik and J. Nakamura, 2010: Mechanisms of tropical  
 1402 Atlantic SST influence on North American precipitation variability. J. Climate, 23, 5610-  
 1403 5628.  
 1404 Lau, K. M., Kim, M. K., and Kim, K. M., 2006: Asian monsoon anomalies induced by  
 1405 aerosol direct effects, Clim. Dynam., 26, 855–864, doi:10.1007/s00382-006-0114-z.  
 1406  
 1407 Li, H., A. Dai, T. Zhou, J. Lu, 2010: Responses of East Asian summer monsoon to  
 1408 historical SST and atmospheric forcing during 1950-2000, Clim. Dyn., 34, 501-514, DOI  
 1409 10.1007/s00382-008-0482-7.  
 1410



1411 Li, J., R. Yu, W. Yuan, H. Chen, W. Sun, Y. Zhang, 2015: Precipitation over East Asia  
 1412 simulated by NCAR CAM5 at different horizontal resolutions. *Journal of Advances in*  
 1413 *Modeling Earth Systems*, 7(2), 774-790. DOI 10.1002/2014MS000414.  
 1414  
 1415 Li, S. L., J. Lu, G. Huang, and K. M. Hu, 2008: Tropical Indian Ocean basin warming  
 1416 and East Asian summer monsoon: A multiple AGCM study. *J. Climate*, 21, 6080–6088.  
 1417  
 1418 Liebmann, B., M.P. Hoerling, C. Funk, I. Bladé, R.M. Dole, D. Allured, X. Quan, P.  
 1419 Pegion, and J.K. Eischeid, 2014: Understanding Recent Eastern Horn of Africa Rainfall  
 1420 Variability and Change. *J. Climate*, 27, 8630–8645. doi: [http://dx.doi.org/10.1175/JCLI-](http://dx.doi.org/10.1175/JCLI-D-13-00714.1)  
 1421 [D-13-00714.1](http://dx.doi.org/10.1175/JCLI-D-13-00714.1)  
 1422  
 1423 Lloyd-Hughes B., J. Hannaford, S. Parry, C. Keef, C. Prudhomme and H.G. Rees, 2010:  
 1424 Drought Catalogues for UK and Europe. Environment Agency Science Report  
 1425 SC070079/SR Environment Agency: Bristol.  
 1426  
 1427 Losada, T., B. Rodriguez-Fonseca, E. Mohino, J. Bader, S. Janicot, and C. R. Mechoso, 2012:  
 1428 Tropical SST and Sahel rainfall: A non-stationary relationship. *Geophys. Res. Lett.*, 39, L12705,  
 1429 doi:[10.1029/ 2012GL052423](https://doi.org/10.1029/2012GL052423).  
 1430  
 1431 Lyon, B. and D.G. DeWitt, 2012: A recent and abrupt decline in the East African long  
 1432 rains. *Geophys. Res. Lett.*, 39, L02702, doi: [10.1029/2011GL050337](https://doi.org/10.1029/2011GL050337).  
 1433

1434 Lyon, B., A.G. Barnston, and D.G. DeWitt, 2013: Tropical Pacific forcing of a 1998-99  
 1435 climate shift: Observational analysis and climate model results for the boreal spring  
 1436 season. *Climate Dyn.*, 43, 893-909, doi 10.1007/s00382-013-1891-9.  
 1437  
 1438 Lyon, B., 2014: Seasonal Drought in the Greater Horn of Africa and its Recent Increase  
 1439 During the March-May Long Rains. *J. Climate*, 27, 7953-7975.  
 1440  
 1441 Ma, H. -Y., C. R. Mechoso, Y. Xue, H. Xiao, C. -M. Wu, J. -L. Li, and F. De Sales,  
 1442 2010: Impact of land surface processes on the South American warm season climate.  
 1443 *Clim. Dyn.* doi: 10.1007/s00382-010-0813-3.  
 1444  
 1445 Magaña, Victor, Jorge A. Amador, and Socorro Medina, 1999: The Midsummer Drought  
 1446 over Mexico and Central America. *J. Climate*, **12**, 1577–1588.  
 1447 doi: [http://dx.doi.org/10.1175/1520-0442\(1999\)012<1577:TMDOMA>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1999)012<1577:TMDOMA>2.0.CO;2)  
 1448  
 1449 Marengo, J. A., and Coauthors, 2008: The drought of Amazonia in 2005. *J. Climate*, 21,  
 1450 495–516.  
 1451  
 1452 Marengo, J. A., R. Jones, L. M. Alves and M. C. Valverde, 2009: Future change of  
 1453 temperature and precipitation extremes in South America as derived from the PRECIS  
 1454 regional climate modeling system. *Int. J. Climatol.*, 29, 2241-2255.  
 1455

1456 Mariotti, A., 2007: How ENSO impacts precipitation in southwest central Asia, *Geophys.*  
1457 *Res. Lett.*, 34, L16706, doi:10.1029/2007GL030078.

1458

1459 Mariotti A., N. Zeng , J. Yoon, V. Artale, A. Navarra, P. Alpert, and L.Z.X. Li, 2008:  
1460 Mediterranean water cycle changes: transition to drier 21st century conditions in  
1461 observations and CMIP3 simulations. *Environ.. Res. Lett.*, 3, 044001 (8pp),  
1462 doi:10.1088/1748-9326/3/4/044001.

1463

1464 Mariotti, A. and A. Dell'Aquila, 2012: Decadal climate variability in the Mediterranean  
1465 region: roles of large-scale forcings and regional processes, *Clim. Dyn.*, 38, 1129–1145,  
1466 DOI 10.1007/s00382-011-1056-7.

1467

1468 Mariotti A., Y. Pan, N. Zeng, and A. Alessandri, 2015: Long-term climate change in the  
1469 Mediterranean region in the midst of decadal variability. *Climate Dyn.*, 44 (5-6), pp  
1470 1437-1456, doi:10.1007/s00382-015-2487-3.

1471

1472 McCarthy et al. (2001): Climate Change 2001: Working Group II: Impacts, Adaptation  
 1473 and Vulnerability. Chapter 14. 14.1.2.1.4. Variability and impacts from El Niño and the  
 1474 Southern Oscillation. [McCarthy, J. J.; Canziani, O. F.; Leary, N. A.; Dokken, D. J.; and  
 1475 White, K. S., ed.], Cambridge University Press, ISBN 0-521-80768-9 (pb: 0-521-01500-  
 1476 6).  
 1477  
 1478 Mechoso, C. R., and S. W. Lyons, 1988: On the atmospheric response to SST anomalies  
 1479 associated with the Atlantic warm event during 1984. J. Climate, 1, 422-428.  
 1480  
 1481 Mechoso, C. R., A. Robertson, and coauthors, 1995: The seasonal cycle over the tropical  
 1482 Pacific in coupled ocean-atmosphere general circulation models. Mon. Wea. Rev., 123,  
 1483 2825–2838.  
 1484  
 1485 Mo, K. C., and E. H. Berbery, 2011: Drought and persistent wet spells over South  
 1486 America based on observations and the U.S. CLIVAR drought experiments. J. Climate,  
 1487 24, 1801-1820.  
 1488  
 1489 Mohino, E., S. Janicot, and J. Bader, 2011a: Sahelian rainfall and decadal to multidecadal  
 1490 SST variability. Climate Dyn., 37, 419–440, doi:[10.1007/s00382-010-0867-2](https://doi.org/10.1007/s00382-010-0867-2).  
 1491  
 1492 Mohino, E., Belén Rodríguez-Fonseca, Teresa Losada, Sébastien Gervois, Serge Janicot,  
 1493 Juergen Bader, Paolo Ruti, Fabrice Chauvin, 2011: [Changes in the interannual SST-forced](#)

1494 [signals on West African rainfall. AGCM intercomparison](#) Climate dynamics 37 (9-10),  
 1495 1707-1725.  
 1496  
 1497 Molod, A., L. Takacs, M. Suarez, J. Bacmeister, I.-S. Song, and A. Eichmann, 2012: The  
 1498 GEOS-5 Atmospheric General Circulation Model: Mean Climate and Development  
 1499 from MERRA to Fortuna. NASA Tech. Rep. Series on Global Modeling and Data  
 1500 Assimilation, NASA TM—2012-104606, Vol. 28, 117 pp.  
 1501  
 1502 Mueller, B., and S.I. Seneviratne, 2012: Hot days induced by precipitation deficits at the  
 1503 global scale. Proc. Natl Acad. Sci., 109 (31), 12398-12403, doi:  
 1504 10.1073/pnas.1204330109  
 1505  
 1506 Müller, O. V., E. H. Berbery, D. Alcaraz-Segura, and M. B. Ek, 2014: Regional Model  
 1507 Simulations of the 2008 Drought in Southern South America Using a Consistent Set of  
 1508 Land Surface Properties. J. Climate, 27, 6754–6778.  
 1509 doi: <http://dx.doi.org/10.1175/JCLI-D-13-00463.1>  
 1510  
 1511 NDMC (2014): ENSO and Drought Around the World.  
 1512 (<http://drought.unl.edu/DroughtBasics/ENSOandForecasting.aspx>).  
 1513  
 1514 Parry, S., J. Hannaford, B. Lloyd-Hughes, C. Prudhomme, and C. Keef, 2010:  
 1515 Drought Summaries of Spatio-temporal evolution of major European Droughts since

1516 1960. Environment Agency Science Report SC070079, Environment Agency, Bristol.  
 1517

1518 Poveda, G. and O. J. Mesa, 1997: Feedbacks between Hydrological Processes in Tropical  
 1519 South America and Large-Scale Ocean–Atmospheric Phenomena. *J. Climate*, 10, 2690–  
 1520 2702.  
 1521 doi: [http://dx.doi.org/10.1175/1520-0442\(1997\)010<2690:FBHPIT>2.0.CO;2](http://dx.doi.org/10.1175/1520-0442(1997)010<2690:FBHPIT>2.0.CO;2)  
 1522

1523 Qian C., and T. Zhou, 2014: Multidecadal variability of North China aridity and its  
 1524 relationship to PDO during 1900-2010, *J. Climate*, 27(3), 1210-1222.  
 1525

1526 Orłowsky, B., and S.I. Seneviratne, 2012: Global changes in extreme events: Regional  
 1527 and seasonal dimension. *Climatic Change*, 110, 669-696, doi: 10.1007/s10584-011-0122-  
 1528 9.  
 1529

1530 Orłowsky, B. and S.I. Seneviratne, 2013: Elusive drought: uncertainty in observed trends  
 1531 and short- and long-term CMIP5 projections, *Hydrol. Earth Syst. Sci.*, 17, 1765-1781,  
 1532 doi:10.5194/hess-17-1765-2013.  
 1533

1534 Orth, R. and S.I. Seneviratne, 2014. Predictability of soil moisture and streamflow on  
 1535 sub-seasonal timescales: A case study. *J. Geophysical Res. - Atmospheres*, 118: 10963-  
 1536 10979.  
 1537

1538 Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J. T., Washington, W.  
 1539 M., Fu, Q., Sikka, D. R., and Wild, M., 2005: Atmospheric brown clouds: Impacts on  
 1540 South Asian climate and hydrological cycle, *Proc. Natl. Acad. Sci. USA*, 102, 5326–  
 1541 5333.  
 1542  
 1543 Rayner, N.A., D.E. Parker, E.B. Horton, C.K. Folland, L.V. Alexander, D.P. Rowell,  
 1544 E.C. Kent, and A. Kaplan, 2003: Global analyses of sea surface temperature, sea ice, and  
 1545 night marine air temperature since the late Nineteenth Century. *J. Geophys. Res.*, 108,  
 1546 (D14), 4407. doi:10.1029/2002JD002670  
 1547  
 1548 Richter, I., 2015: Climate model biases in the eastern tropical oceans: causes, impacts and  
 1549 ways forward. *Wiley Inter. Rev: Clim. Change*, 6, 345–358, doi:10.1002/wcc.338.  
 1550  
 1551 Rienecker, M. M., and Coauthors, 2008: The GEOS-5 data assimilation system—  
 1552 Documentation of versions 5.0.1, 5.1.0, and 5.2.0. NASA Tech. Rep. Series on Global  
 1553 Modeling and Data Assimilation, NASA/TM-2007-104606, Vol. 27, 95 pp.  
 1554  
 1555 Robson, J., R. Sutton, K. Lohmann, D. Smith, M. Palmer, 2012: The causes of the rapid  
 1556 warming of the North Atlantic Ocean in the mid 1990s. *J. Climate*, 25, 4116-4134.  
 1557  
 1558 Rodell, M., P. R. Houser, U. Jambor, J. Gottschalck, K. Mitchell, C.-J. Meng, K.  
 1559 Arsenault, B. Cosgrove, J. Radakovich, M. Bosilovich, J. K. Entin, J. P. Walker, D.  
 1560 Lohmann, and D. Toll, The Global Land Data Assimilation System, *Bull. Amer. Meteor.*

1561 Soc., 85(3): 381-394, 2004.

1562

1563 Rodríguez-Fonseca, B., I. Polo, J. García-Serrano, T. Losada, E. Mohino, C. R. Mechoso,  
 1564 and F. Kucharski, 2009: Are Atlantic Niños enhancing Pacific ENSO events in recent  
 1565 decades? *Geophys. Res. Lett.*, 36, L20705, doi:10.1029/2009GL040048.

1566

1567 Rodríguez-Fonseca, B., S. Janicot, E. Mohino, T. Losada, J. Bader, , Cyril Caminade,  
 1568 Fabrice Chauvin, Bernard Fontaine, Javier García-Serrano, Sebastien Gervois, Mathieu  
 1569 Joly, Irene Polo, Paolo Ruti, Pascal Roucou, Aurore Voldoire, 2011:  
 1570 Interannual and decadal SST-forced responses of the West African monsoon.  
 1571 *Atmospheric Science Letters*, 12(1), 67-74.

1572

1573 Rodríguez-Fonseca, B., E. Mohino, C.R. Mechoso, C. Caminade, M. Biasutti, M.  
 1574 Gaetani, J. Garcia-Serrano, E.K. Vizy, K. Cook, Y. Xue, I. Polo, T. Losada, L. Druyan,  
 1575 B. Fontaine, J. Bader, F.J. Doblas-Reyes, L. Goddard, S. Janicot, A. Arribas, W. Lau, A.  
 1576 Colman, M. Vellinga, D.P. Rowell, F. Kucharski, and A. Voldoire, 2015: Variability and  
 1577 predictability of West African droughts: A review of the role of sea surface temperature  
 1578 anomalies. *J. Climate*, in press, doi:10.1175/JCLI-D-14-00130.

1579

1580 Roeckner, E., G. Bäuml, L. Bonaventura, R. Brokopf, M. Esch, M. Giorgetta, S.  
 1581 Hagemann, I. Kirchner, L. Kornblueh, E. Manzini, A. Rhodin, U. Schlese, U.  
 1582 Schulzweida, and A. Tompkins, 2003: The atmospheric general circulation model  
 1583 ECHAM5. Part I: Model description. Max Planck Institute for Meteorology Rep. 349,



1584 127 pp. [available from MPI for Meteorology, Bundesstr. 53, 20146 Hamburg,  
 1585 Germany].

1586

1587 Ropelewski, C. F., and M. S. Halpert ,1987: Global and regional scale precipitation  
 1588 patterns associated with the El Niño/ Southern Oscillation. Mon. Wea. Rev., 115, 1606-  
 1589 1626.

1590

1591 Rowell, D. P., 2013: Simulating SST Teleconnections to Africa: What is the State of the  
 1592 Art?. J. Climate, 26, 5397–5418. doi: <http://dx.doi.org/10.1175/JCLI-D-12-00761.1>

1593

1594 [Sánchez, E. and coauthors, 2015: Regional climate modelling in CLARISLPB: a](#)  
 1595 [concerted approach towards twenty-first century projections of regional temperature and](#)  
 1596 [precipitation over South America, Climate Dyn. DOI 10.1007/s00382-014-2466-0.](#)

1597

1598 Saravanan R. and P. Chang, 2000: Interaction between tropical Atlantic variability and  
 1599 El Niño-Southern Oscillation. J. Climate 13, 2177–2194.

1600

1601 Scaife, A.A., J.R. Knight, G.K. Vallis, C.K. Folland, 2005: A stratospheric influence on  
 1602 the winter NAO and North Atlantic surface climate. Geophys. Res. Lett. 32, L18715.

1603

1604 Schubert, S.D., M. J. Suarez, P. J. Pegion, R. D. Koster, and J. T. Bacmeister , 2004: On  
 1605 the Cause of the 1930s Dust Bowl, Science, **33**, 1855-1859.

1606 Schubert, S., D. Gutzler, H. Wang, A. Dai, T. Delworth, C. Deser, K. Findell, R. Fu, W.  
 1607 Higgins, M. Hoerling, B. Kirtman, R. Koster, A. Kumar, D. Legler, D. Lettenmaier, B.  
 1608 Lyon, V. Magana, K. Mo, S. Nigam, P. Pegion, A. Phillips, R. Pulwarty, D. Rind, A.  
 1609 Ruiz-Barradas, J. Schemm, R. Seager, R. Stewart, M. Suarez, J. Syktus, M. Ting, C.  
 1610 Wang, S. Weaver, N. Zeng, 2009: A USCLIVAR Project to Assess and Compare the  
 1611 Responses of Global Climate Models to Drought-Related SST Forcing Patterns:  
 1612 Overview and Results. *J. Climate*, 22, 5251–5272.  
 1613  
 1614 Schubert, S., H. Wang, R. Koster, M. Suarez, and P. Groisman, 2014: Northern Eurasian  
 1615 Heat Waves and Droughts. *J. Climate*, 27, 3169-3207.  
 1616  
 1617 Seager, R., Y. Kushnir, C. Herweijer, N. Naik, and J. Velez, 2005: Modeling the tropical  
 1618 forcing of persistent droughts and pluvials over western North America: 1856-2000. *J.*  
 1619 *Climate*, 18, 4068-4091.  
 1620  
 1621 Seager, R., N. Naik, W. Baethgen, A. Robertson. Y. Kushnir, J. Nakamura, S. Jurburg,  
 1622 2010: Tropical oceanic causes of interannual to multidecadal precipitation variability in  
 1623 southeast South America over the past century. *J. Climate*, 23, 5517-5539.  
 1624  
 1625 Seager, R., L. Goddard, J. Nakamura, N. Henderson, D. E. Lee, 2014a: Dynamical causes  
 1626 of the 2010/11 Texas-northern Mexico drought. *J. Hydrometeor.* 2013, 15, 39-68.  
 1627

1628 Seager, R., H. Liu, N. Henderson, I. Simpson, C. Kelley, T. Shaw, Y. Kushnir and M.  
 1629 Ting, 2014b: Causes of increasing aridification of the Mediterranean region in response  
 1630 to rising greenhouse gases, *J. Climate*, 27, 4655-4676.  
 1631  
 1632 Seager, R., M. Hoerling, S. Schubert, H. Wang, B. Lyon, A. Kumar, J. Nakamura, and N.  
 1633 Henderson, 2014c: Causes and predictability of the 2011–14 California drought,  
 1634 DTF/NIDIS Assessment Report, 40 pp., doi:10.7289/V7258K7771F.  
 1635  
 1636 Seager, R. and M. Hoerling, 2014: Atmosphere and Ocean Origins of North American  
 1637 Drought. *J. Climate*, 27, 4581-4606.  
 1638  
 1639 Seneviratne, S. I. and Coauthors, 2006: Soil moisture memory in AGCM simulations:  
 1640 Analysis of Global Land–Atmosphere Coupling Experiment (GLACE) data. *J.*  
 1641 *Hydrometeor.*, 7, 1090–1112.  
 1642  
 1643 Seneviratne, S.I., N. Nicholls, D. Easterling, C.M. Goodess, S. Kanae, J. Kossin, Y. Luo,  
 1644 J. Marengo, K. McInnes, M. Rahimi, M. Reichstein, A. Sorteberg, C. Vera, and X.  
 1645 Zhang, 2012a: Changes in climate extremes and their impacts on the natural physical  
 1646 environment. In: *Managing the Risks of Extreme Events and Disasters to Advance*  
 1647 *Climate Change Adaptation* [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken,  
 1648 K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.-K. Plattner, S.K. Allen, M. Tignor, and P.M.  
 1649 Midgley (eds.)]. A Special Report of Working Groups I and II of the Intergovernmental  
 1650 Panel on Climate Change, pp. 109-230.

1651

1652 Seneviratne, S.I., I. Lehner, J. Gurtz, A.J. Teuling, H. Lang, U. Moser, D. Grebner, L.  
1653 Menzel, K. Schroff, T. Vitvar, and M. Zappa, 2012b: Swiss pre-alpine Rietholzbach  
1654 research catchment and lysimeter: Analysis of 32--year hydroclimatological time series  
1655 and 2003 drought. *Water Resources Research*, 48, W06526,  
1656 doi:10.1029/2011WR011749.

1657

1658 Sheffield, J., E.F. Wood, and M.L. Roderick, 2012: Little change in global drought in the  
1659 past 60 years. *Nature*, 491, 435-438, doi:10.1038/nature11575.

1660

1661 Smith, D.M., A.A. Scaife, and B. P. Kirtman, 2012: What is the current state of  
1662 knowledge with regard to seasonal to decadal forecasting? *Environ. Res. Lett.*, 7, 015602  
1663 (11pp).

1664

1665 Song F. and T. Zhou, 2014a: Interannual Variability of East Asian Summer Monsoon  
1666 Simulated by CMIP3 and CMIP5 AGCMs: Skill Dependence on Indian Ocean–Western  
1667 Pacific Anticyclone Teleconnection. *J. Climate*, 27, 1679–1697.

1668

1669 Song F., and T. Zhou, 2014b: The climatology and interannual variability of East Asian  
1670 summer monsoon in CMIP5 coupled models: Does air-sea coupling improve the  
1671 simulations? *J. Climate*, 27, 8761-8777

1672

1673 Song F., T. Zhou, and Y. Qian, 2014: Responses of East Asian summer monsoon to  
 1674 natural and anthropogenic forcings in the 17 latest CMIP5 models, *Geophys. Res. Lett.*,  
 1675 41, doi:10.1002/2013GL058705  
 1676  
 1677 Sörensson, A. A., and C. G. Menéndez, 2011: Summer soil–precipitation coupling in  
 1678 South America. *Tellus A*, 63, 56–68. doi: 10.1111/j.1600-0870.2010.00468.x  
 1679  
 1680 Spinoni, J., G. Naumann, J. Vogt, and P. Barbosa, 2015: The biggest drought events in  
 1681 Europe from 1950 to 2012. *J. Hydrology*, in press.  
 1682  
 1683 Stagge, J.H., I. Kohn, L.M. Tallaksen, and K. Stahl, 2015: Modeling drought impact  
 1684 occurrence based on meteorological drought indices in Europe. *J. Hydrology*, 530, 37-50.  
 1685  
 1686 Stahl, K., 2001: Hydrological Drought – a Study across Europe. PhD Thesis, Albert-  
 1687 Ludwigs-Universität Freiburg, Freiburger Schriften zur Hydrologie no. 15, also available  
 1688 from: <<http://www.freidok.unifreiburg.de/volltexte/202>>.  
 1689  
 1690 Staudinger, M., K. Stahl, and J. Seibert, 2014: A drought index accounting for snow.  
 1691 *Water Resour. Res.*, 50, 7861–7872, doi:10.1002/2013WR015143.  
 1692  
 1693 Sutton, R. and B. Dong, 2012: Atlantic Ocean influence on a shift in European climate in  
 1694 the 1990s. *Nature Geoscience*, 5, pp. 788-792. ISSN 1752-0908 doi: 10.1038/ngeo1595.  
 1695

1696 Tallaksen, L.M., J.H. Stagge, K. Stahl, L. Gudmundsson, R. Orth, S.I. Seneviratne, A.F.  
 1697 van Loon, and H.A.J. van Lanen, 2015: Characteristics and drivers of drought in Europe  
 1698 – a summary of the DROUGHT-RSPI project. In: “Drought: Research and Science-  
 1699 Policy Interfacing”, Andreu et al. (eds). Taylor & Francis Group, London, ISBN 978-1-  
 1700 138-02779-4.  
 1701  
 1702 Teuling, A.J., A. van Loon, S.I. Seneviratne, I. Lehner, M. Aubinet, B. Heinesch, C.  
 1703 Bernhofer, T. Grünwald, H. Prasse, and U. Spank, 2013: Evapotranspiration amplifies  
 1704 European summer drought. *Geophys. Res. Lett.*, 40 (10), 2071-2075  
 1705  
 1706 Van Loon, A. F., and H.A. Van Lanen, 2012. A process-based typology of hydrological  
 1707 drought, *Hydrol. Earth Syst. Sci.*, 16: 1915–1946.  
 1708  
 1709 Vicente-Serrano, Sergio M., Santiago Beguería, and Juan I. López-Moreno, 2010: A  
 1710 Multiscalar Drought Index Sensitive to Global Warming: The Standardized Precipitation  
 1711 Evapotranspiration Index. *J. Climate*, **23**, 1696–1718.  
 1712 doi: <http://dx.doi.org/10.1175/2009JCLI2909.1>  
 1713  
 1714 Wang, G., Y. Kim, and D. Wang, 2007: Quantifying the strength of soil moisture-  
 1715 precipitation coupling and its sensitivity to changes in surface water budget. *J.*  
 1716 *Hydrometeorol.*, 8, 551–570.  
 1717

1718 Wang, C., S.-K. Lee, and D. B. Enfield, 2008: Climate Response to Anomalously Large  
 1719 and Small Atlantic Warm Pools during the Summer. *J. Climate*, 21, 2437–2450. doi:  
 1720 <http://dx.doi.org/10.1175/2007JCLI2029.1>  
 1721  
 1722 Wang, H., S.D. Schubert, M. J. Suarez, J. Chen, M. Hoerling, A. Kumar and P. Pegion,  
 1723 2009: Attribution of the seasonality and regionality in climate trends over the United  
 1724 States during 1950-2000. *J. Climate*, 2571-2590.  
 1725  
 1726 Wang, H., S. Schubert, R. Koster, Y.-G. Ham, and M. Suarez, 2014: On the Role of SST  
 1727 Forcing in the 2011 and 2012 Extreme U.S. Heat and Drought: A Study in Contrasts. *J.*  
 1728 *Hydrometeor*, **15**, 1255–1273.  
 1729 doi: <http://dx.doi.org/10.1175/JHM-D-13-069.1>  
 1730  
 1731 Wu B., T. Zhou, T. Li, 2009, Seasonally Evolving Dominant Interannual Variability  
 1732 Modes of East Asian Climate, *J. Climate*, 22, 2992-3005  
 1733  
 1734 Wu B., T. Li, and T. Zhou, 2010: Relative contributions of the Indian Ocean and local  
 1735 SST anomalies to the maintenance of the western North Pacific anomalous anticyclone  
 1736 during El Nino decaying summer. *J. Climate*, 23, 2974-2986.  
 1737  
 1738 Wu R., Z. Hu, and B. Kirtman, 2003: Evolution of ENSO-Related Rainfall Anomalies in  
 1739 East Asia. *J. Climate*, 16, 3742–3758.  
 1740

1741

1742 Xie, S.-P., K. Hu, J. Hafner, H. Tokinaga, Y. Du, G. Huang, and T. Sampe, 2009: Indian  
1743 Ocean capacitor effect on Indowestern Pacific climate during the summer following El  
1744 Niño. *J. Climate*, 22, 730–747.

1745

1746 Xoplaki, E., J. F. Gonzalez-Rouco and J. Luterbacher, 2004: Wet season Mediterranean  
1747 precipitation variability: influence of large-scale dynamics and trends. *Clim. Dyn.*, 23,  
1748 63-78.

1749

1750 Xue, Y., F. de Sales, C. R. Mechoso, C. A. Nobre, H. -M. Juang, 2006: Role of Land  
1751 Surface Processes in South American Monsoon Development. *J. Climate*, 19, 741-762.

1752

1753 Yang, W., R. Seager, M.A. Cane, and B. Lyon, 2014: The East African Long Rains in  
1754 Observations and Models. *J. Climate*, 27, 7185-7202.

1755

1756 Yang, F. L., and K. M. Lau (2004), Trend and variability of China precipitation in spring  
1757 and summer: Linkage to sea-surface temperatures, *Int. J. Climatol.*, 24, 1625 – 1644,  
1758 doi:10.1002/joc.1094.

1759

1760 Yin, D., M.L. Roderick, G. Leech, F. Sun, and Y. Huang, 2014: The contribution of  
1761 reduction in evaporative cooling to higher surface air temperatures during drought.  
1762 *Geophys. Res. Lett.*, doi:10.1002/2014GL062039.

1763



1764 Yoon, J.-H., and N. Zeng, 2009: An Atlantic influence on Amazon rainfall. *Climate Dyn.*,  
 1765 34, 249–264.  
 1766  
 1767 Zeng, N., J. D. Neelin, K.-M. Lau, and C. J. Tucker, 1999: Enhancement of interdecadal  
 1768 climate variability in the Sahel by vegetation interaction. *Science*, 286, 1537-1540.  
 1769  
 1770 Zhang, L. and T. Zhou, 2015: Drought over East Asia: A Review, *J. Climate*, 28(8),  
 1771 3375-3399.  
 1772  
 1773 Zhou T., R. Yu, H. Li, and B. Wang, 2008a: Ocean Forcing to Changes in Global  
 1774 Monsoon Precipitation over the Recent Half-Century, *J. Climate*, 21(15), 3833–3852.  
 1775  
 1776 Zhou, T., L. Zhang, and H. Li, 2008b: Changes in global land monsoon area and total  
 1777 rainfall accumulation over the last half century, *Geophys. Res. Lett.*,  
 1778 doi:10.1029/2008GL034881.  
 1779  
 1780 Zhou, T., D. Gong, J. Li, and B. Li, 2009a: Detecting and understanding the multi-  
 1781 decadal variability of the East Asian Summer Monsoon – Recent progress and state of  
 1782 affairs. *Meteorologische Zeitschrift*, 18 (4), 455-467.  
 1783  
 1784 Zhou T., Wu B., B. Wang, 2009b: How well do atmospheric general circulation models  
 1785 capture the leading modes of the interannual variability of the Asian-Australian  
 1786 Monsoon?, *J. Climate*, 22, 1159-1173.

1787

1788 Zhou, T., Bo Wu, A. A. Scaife, S. Bronnimann, A. Cherchi, D. Fereday, A. M. Fischer, C.

1789 K. Folland, K.E. Jin, J. Kinter, J. R. Knight, F. Kucharski, S. Kusunoki, N.-C. Lau,

1790 Lijuan Li, M.J. Nath, T. Nakaegawa, A. Navarra, P. Pegion, E. Rozanov, S. Schubert, P.

1791 Sporyshev, A. Voldoire, Xinyu Wen, J. H. Yoon ,N. Zeng, 2009c: The CLIVAR C20C

1792 Project: Which components of the Asian-Australian Monsoon circulation variations are

1793 forced and reproducible? *Climate Dyn.*, 33, 1051–1068.

1794

1795 Zhou, T., F. Song, R. Lin, X. Chen and X. Chen, 2013: The 2012 North China floods:

1796 Explaining an extreme rainfall event in the context of a long-term drying tendency [in

1797 “Explaining Extreme Events of 2012 from a Climate Perspective”]. *Bull. Amer. Meteor.*

1798 *Soc.*, 94(9), S49-S51.

1799

1800 Zwiers, F. W., X. L. Wang and J. Sheng, 2000: Effects of specifying bottom boundary


1801 conditions in an ensemble of atmospheric GCM simulations. *J. Geophys. Res.*, 105, 7295-7315.

1802

*Table 1: Temporal correlation between observed annual mean regional mean precipitation (GPCP) and 1) the regional mean annual mean PDSI from Dai et al. (2004) for 1979-2005, and 2) the regional mean annual mean soil moisture (top 100cm) from the Global Land Data Assimilation System Version 2 (GLDAS-2, Rodell et al. 2004) for 1979-2010. The numbers (1-10) in the Table refer to the regions outlined in Figures 5-8 as follows: Figure 5: 1 (US and N. Mexico), 2 (N. South America and Central America), 3 (central South America). Figure 6: 4 (west Africa), 5 (east Africa). Figure 7: 6 (Mideast), 7 (southern Asia), 8 (east Asia). Figure 8: 9 (Australia), 10 (Indonesia).*

	1	2	3	4	5	6	7	8	9	10
PDSI (1979-2005)	0.52	0.80	0.69	0.66	0.66	0.72	0.69	0.65	0.75	0.71
Soil Moisture (1979-2010)	0.72	0.86	0.76	0.70	0.57	0.80	0.66	0.50	0.80	0.81

1813



1814

1815 *Figure 1. (Top) The background map shows the ratio of two variances: the variance of the*  
 1816 *ensemble mean time series of annual precipitation and the total variance of annual mean*  
 1817 *precipitation over all ensemble members (ratios are computed for each model separately and*  
 1818 *then averaged). Higher values of the ratio indicate a stronger impact of the prescribed SSTs on*  
 1819 *the precipitation time series. The small maps show the correlations between the ensemble mean*  
 1820 *annual fields (averaged over the boxed areas) with SST (correlations are computed for each*  
 1821 *model separately and then averaged). Results are based on 12 ensemble members for each of 5*

1822 *models (GEOS-5, CCM3, CAM4, GFS, ECHAM5) using detrended values for the period 1979-*  
1823 *2011. (Bottom) Same, but for 2m air temperature (note change in contour interval). The*  
1824 *horizontal color bars are for the variance ratios, and the vertical color bars are for the*  
1825 *correlations.*  
1826





Figure 2: The shift in probabilities of extremes between the two periods 1998-2011 and 1979-1993 defined as  $(P(x_2 > x_c) - P(x_1 > x_c)) / P(x > x_c)$ , where  $x_2$  refers to values during the recent period (1998-2011) and  $x_1$  refers to values during the earlier period (1979-1993). The shift is normalized by  $P(x > x_c)$ , where  $x$  refers to values during the entire time period,

1832 and  $x_c$  is chosen so that  $P(x > x_c)$  is 2.5%. The top panel shows results for precipitation  
 1833 and the bottom for 2m temperature; in the case of precipitation, the shift in probability  
 1834 actually refers to the left tail of the distribution (values less than  $x_c$ ). The results are  
 1835 based on 12 ensemble members for each of 5 models (GEOS-5, CCM3, CAM4, GFS,  
 1836 ECHAM5). Each model's values are first normalized to have zero mean and unit  
 1837 variance. The inserts show the actual PDFs for the two periods (red is for the recent  
 1838 period and blue indicates the earlier period) for all grid points in the indicated boxes,  
 1839 land only). Vertical lines highlight the zero value and the value of  $x_c$ .  
 1840



1841  
 1842 Figure 3: Top left: Mean simulated precipitation differences between 1998-2011 and 1979-1993,  
 1843 based on results from the five models. Bottom left: Corresponding differences in T2M (land  
 1844 only). Middle panels: Same as left panels, but for the observations. Top right: The mean  
 1845 observed SST differences between 1998-2011 and 1979-1993. Bottom right: the time series of the  
 1846 PDO and AMO.




Figure 4: Observed (GPCP, solid lines) and simulated (5-model ensemble mean, dashed lines) annual cycle of precipitation (mm/day) for the selected regions based on the period 1979-2011. The regions (land only) are those examined in Figure 5, Figure 6, Figure 7, and Figure 8 (see the boxes outlining the regions in those figures).






Figure 5: Left panels: The correlations between the ensemble mean precipitation averaged over the United States and northern Mexico (black box) and SST for individual seasons (correlations are averaged over the 5 models). Middle panels: same as left panels, but for northern South America and Central America (black box). Right panels: Same as left panels, but for central South America (black box).




Figure 6: Left panels: The correlations between the ensemble mean precipitation averaged over West Africa (black box) and SST for individual seasons ((correlations are averaged over the 5 models)). Right panels: Same, but for east Africa.




Figure 7: Left panels: The correlations between the ensemble mean precipitation averaged over the middle east (black box) and SST for individual seasons (correlations are averaged over the 5 models). Middle panels: Same, but for southern Eurasia. Right panels: Same, but for east Asia.




Figure 8: Left panels: Correlations between the ensemble mean precipitation averaged over Australia (black box) and SST for individual seasons (correlations are averaged over the 5 models). Right panels: Same, but for Indonesia.







Figure 9: The 5-model mean simulated precipitation differences between 1998-2011 and 1979-1993 for individual months. Units: mm/day.




1880

1881 *Figure 10: Same as Figure 2a, but for each season.*

1882

1883



1884

1885 *Figure 11: Same as Figure 2b, but for each season.*

1886






Figure B1: The correlations between the ensemble mean precipitation over East Africa and SST for each model during SON (1979-2011). Also shown (top middle panel) is the correlation based on GPCP observations.






Figure B2: The correlations between the ensemble mean annual precipitation over the Middle East and SST for each model (1979-2011). Also shown (top middle panel) is the correlation based on GPCP observations. We note that there is some sensitivity to the



1896 region chosen for some of the models (e.g. the results for CCM3 looks more like that of  
 1897 the other models and observations if the region is truncated on the southern and eastern  
 1898 edges).




1899  
 1900 *Figure B3: The correlations between the ensemble mean precipitation over southern Asia*  
 1901 *and SST for each model for MAM (1979-2011). Also shown (top middle panel) is the*  
 1902 *correlation based on GPCP observations.*



1904

1905 *Figure B4: The correlations between the ensemble mean precipitation over Australia and*  
1906 *SST for each model for SON (1979-2011). Also shown (top middle panel) is the*  
1907 *correlation based on GPCP observations.*



1908  
1909 *Figure B5: The correlations between the ensemble mean precipitation over the southern*  
1910 *US and SST for each model for MAM (1979-2011). Also shown (top middle panel) is the*  
1911 *correlation based on GPCP observations.*

1912

## List of Figures

Figure 1. (Top) The background map shows the ratio of two variances: the variance of the ensemble mean time series of annual precipitation and the total variance of annual mean precipitation over all ensemble members. Higher values of the ratio indicate a stronger impact of the prescribed SSTs on the precipitation time series. The small maps show the correlations between the ensemble mean annual fields (averaged over the boxed areas) with SST. All results are for the period 1979-2011 and are based on 60 ensemble members: 12 AMIP simulations for each of 5 models (GEOS-5, CCM3, CAM4, GFS, ECHAM5). Results are based on detrended values. (Bottom) Same, but for 2m air temperature (note change in contour interval). The horizontal color bars are for the variance ratios, and the vertical color bars are for the correlations.

Figure 2: The shift in probabilities of extremes between the two periods 1998-2011 and 1979-1993 defined as  $(P(x_2 > x_c) - P(x_1 > x_c)) / P(x > x_c)$ , where  $x_2$  refers to values during the recent period (1998-2011) and  $x_1$  refers to values during the earlier period (1979-1993). The shift is normalized by  $P(x > x_c)$ , where  $x$  refers to values during the entire time period, and  $x_c$  is chosen so that  $P(x > x_c)$  is 2.5%. The top panel shows results for precipitation and the bottom for 2m temperature; in the case of precipitation, the shift in probability actually refers to the left tail of the distribution (values less than  $x_c$ ). The results are based on 12 ensemble members for each of 5 models (GEOS-5, CCM3, CAM4, GFS, ECHAM5). Each model's values are first normalized to have zero mean and unit variance. The inserts show the actual PDFs for the two periods (red is for the recent

1935 period and blue indicates the earlier period) for all grid points in the indicated boxes, land  
1936 only). Vertical lines highlight the zero value and the value of  $x_c$ .

1937 Figure 3: Top left: Mean simulated precipitation differences between 1998-2011 and  
1938 1979-1993, based on results from the five models. Bottom left: Corresponding  
1939 differences in T2M (land only). Middle panels: Same as left panels, but for the  
1940 observations. Top right: The mean observed SST differences between 1998-2011 and  
1941 1979-1993. Bottom right: the time series of the PDO and AMO.

1942 Figure 4: Observed (GPCP, solid lines) and simulated (5-model ensemble mean, dashed  
1943 lines) annual cycle of precipitation (mm/day) for the selected regions based on the period  
1944 1979-2011. The regions are those examined in Figure 5, Figure 6, Figure 7, and Figure 8  
1945 (see the boxes outlining the regions in those figures).

1946 Figure 5: Left panels: The correlations between the ensemble mean precipitation  
1947 averaged over the United States and northern Mexico (black box) and SST for individual  
1948 seasons (correlations are averaged over the 5 models). Middle panels: same as left  
1949 panels, but for northern South America and Central America (black box). Right panels:  
1950 Same as left panels, but for central South America (black box).

1951

1952

1953 Figure 6: Left panels: The correlations between the ensemble mean precipitation  
1954 averaged over the Sahel (black box) and SST for individual seasons (5-model mean).

1955 Right panels: Same, but for east Africa.

1956

1957 Figure 7: Left panels: The correlations between the ensemble mean precipitation  
 1958 averaged over the middle east (black box) and SST for individual seasons (5-model  
 1959 mean). Middle panels: Same, but for southern Eurasia. Right panels: Same, but for east  
 1960 Asia.  
 1961  
 1962 Figure 8: Left panels: Correlations between the ensemble mean precipitation averaged  
 1963 over Australia (black box) and SST for individual seasons (5-model mean). Right panels:  
 1964 Same, but for Indonesia.  
 1965  
 1966 Figure 9: The 5-model mean simulated precipitation differences between 1998-2011 and  
 1967 1979-1993 for individual months. Units: mm/day.  
 1968  
 1969 Figure 10: Same as Figure 2a, but for each season.  
 1970  
 1971 Figure 11: Same as Figure 2b, but for each season.  
 1972  
 1973  
 1974 Figure B1: The correlations between the ensemble mean precipitation over East Africa  
 1975 and SST for each model during SON (1979-2011). Also shown (top middle panel) is the  
 1976 correlation based on GPCP observations.  
 1977  
 1978 Figure B2: The correlations between the ensemble mean annual precipitation over the  
 1979 Middle East and SST for each model (1979-2011). Also shown (top middle panel) is the

correlation based on GPCP observations. We note that there is some sensitivity to the region chosen for some of the models (e.g. the results for CCM3 looks more like that of the other models and observations if the region is truncated on the southern and eastern edges).

Figure B3: The correlations between the ensemble mean precipitation over southern Asia and SST for each model for MAM (1979-2011). Also shown (top middle panel) is the correlation based on GPCP observations.

Figure B4: The correlations between the ensemble mean precipitation over Australia and SST for each model for SON (1979-2011). Also shown (top middle panel) is the correlation based on GPCP observations.

Figure B5: The correlations between the ensemble mean precipitation over the southern US and SST for each model for MAM (1979-2011). Also shown (top middle panel) is the correlation based on GPCP observations.